INDEX FOR VOLUME 29

New species in **blackface type**
Joint authorship indicated by pages in ( )
Prepared by Frederick V. Rand, Office of Experiment Stations, U. S. Department of Agriculture

Acer, bacterial leaf spot of spp., 968
blight of spp., pythiaceous fungus associated, 11
butt and trunk rot due to Daedalia, 3
cankers, Nectria, avenues of entry, 353
Polyporus, 3
Aceratagallia sanguinolenta, effect on virus of potato yellow dwarf, 2
Acetic acid, -mercuric chloride-alcohol, for bean bacterial blight seed treatment, 19
in spray residue removal, 12
Actinomyces scabies, as influenced by soil pH, 750
isolation method, 1000
pathogenic races, relation to resistance, 204
potato infection controlled by calomel, 13
Actinomycetes, phytopathogenic, isolation methods, 1000
sweet potato pathogenicity of strains, 17
Aedes aegypti, virus activity inhibited by juices of, 324
Aegilops ventricosa, Tilletia spp. infecting, 576
Agaricus, antagonism between Fusarium and, 728
Agropyron, Tilletias infecting spp. of, 576
Ustilago hordei on spp., 490
trichophorum, Puccinia rubigo-vera tritici infection in, 965
Agrostis stolonifera, degeneration deficiency disease (new) of Metropolitan var., 95
Air, wheat leaf-rust spores disseminated by, 4
Airplane, spraying by, in peach orchards, 823
Albugo occidentalis, spinach rust, 12, 616
epidemic, 12
taxonomic comparisons, 616
Alcohol, -mercuric chloride-acetic acid, for bean bacterial blight seed treatment, 19
Alfalfa, mild mosaic of broad bean infecting, 452
susceptibility to Macrosporium sarcinaeforme vs. Pleospora lycopersici, 541
ALLISON, J. LEWIS, 1, 554, (978)
Allium cepa (See Onion)
Almond, Sclerotinia laxa infection in Calif., 349
Alternaria, hardwoods stained by spp., 1037
pines infected by spp., 1036
solani, tomato blight, control by cuprous oxides, 315
early spraying for, 11
Aluminum sulphate, narcissus bulb rot control by, 23
Amaranthus retroflexus, Corticium microsclerotia on, 560
potato virus-X host, 57
Ambrosia elatior, Corticium microsclerotia on, 560
American Phytopathological Society, Report, Annual Meeting, 375
Summer Meeting, 1003
Ammonium sulphate, in purification of tobacco mosaic virus protein, 251
Amphorophora rubi, raspberries resistant to, breeding and factors in, 647
as vector of raspberry mosaic, 647
rubicumberlandi, as vector of black-raspberry mosaic, 825
Analysis of variance, review, 657
Ananas comosus (See Pineapple)
ANDERSON, E. J., 1
ANDERSON, OLOF CAMPBELL, 26
ANDES, J. O., 750
ANDRUS, C. F., 1, 9
Angular leaf spot, cotton, persistence and distribution factors under irrigation, 754
seed treatment by H_2SO_4 for, 879
tobacco, as influenced by water-soaking, 2, 889
Antagonisms, Agaricus/Fusarium, 728
in Penicillium spp., 235
Anthracnose, bean, resistance to, factorial interpretation, 1
of Lippia lanceolata, Sphaceloma lippiae causing, 654
Antibody, -like substance, to curly top virus, 828
Antirrhinum, rust, resistant lines, 18
majus, Botrytis blight, trichomes as infection courts, 651
rust, infection in relation to moisture, 933
Aphids as vectors, black-raspberry mosaic, 825
broad-bean mild mosaic, 448
celery virus diseases, 824
Aphis rumicis, vector of mild mosaic of broad bean, 450
virus activity inhibited by juices of, 324
Aplanobacter stewarti, Tripsacum dactyloides infected spontaneously by, 6, 284
APPEL, O., 558
Apple, copper fungicide compatibility with arsenical and nicotine-oil sprays, 12
Corticium white root rot, 5
crown gall, effect on young trees, 21
hairy root, effect on young trees, 21
internal bark necrosis, as boron deficiency disease, 10
leaf hopper, Entomophthora sp. from, 9
lime-sulphur causing yield reduction, 6
perennial canker, as influenced by wound dressings, 739
Phytophthora cactorum from Lychnis non-pathogenic on, 654
rusts, pathogenicity, symptoms and 3 fungi compared, 801
varietal susceptibility in relation to spp. of, 18

scab, control by phenothiazine and derivatives, 7, 707
spray residue removal, 12
Arceuthobium, biological control of spp. by fungus infection, 995
campylopodum, stem disease due to Fusarium(?), 995
douglasii, stem disease due to Fusarium(?), 995
ARK, P. A., 633, 968
Armillaria mellea, in orchard soils, carbon bisulphide control, 827
ARMSTRONG, G. M., 1, (23)
Arsenicals, compatibility with copper fungicides, 12
removal from apples, 12
Arthrobotrys musiformis, nematode injury restricted by, 599–,
oligospora, nematode injury restricted by, 600–,
Ascochyta caulicola, sweetclover seed infected by, 912
Ascospores, Sclerotinia, method for forced ejection, 373
Ash, black (See Fraxinus nigra)
white (See Fraxinus americana)
Asimina triloba, virus-like symptoms on, 8
Aspen (See Populus)
Aspergillus, incidence of spp. on cotton, 754
Aster yellows, in delphinium, incubation period and leaf-hopper transmission, 826
relation to potato purple-top, 14
Atropa belladonna, Solanum virus-4 in, 171
Aucuba mosaic, potato, differentiation and classification, 6, 918
Authors, advice to, 292
Autolysis, in Sphaerotheca sorghi crosses, 14
Auxins (See also Growth substances)
in root nodules of Phaseolus, increase in, 15
Avocado, sun-blotch, in Florida, 537
Axonopus compressus, leaf blotch (Ciboria associated), 749
furcatus, leaf blotch (Ciboria associated), 749
AYERS, T. T., (15)
Azalea, bud and twig blight due to Sporocybe azaleae, 517

Bacillus (See also Erwinia)
carotovorus, strains compared, 831
phytophthorus, strains compared, 831
radiobacter, growth substance production on peptone broth by, 16
tracheiphilus, cucumber hybrids resistant to, 996
Bacteria, Dieffenbachia leafspot due to, 19, 956
infection by, as influenced by water-soaking, 2
as influenced by nitrogen supply, 1051
lespedeza wilt due to possible **n. sp.**, 15
motile, corn wilt-like symptoms induced by, 1062
phytopathogenic, poured-plate vs. single-cell methods for studying, 852
taxonomy, 128
transfer hood for cultures, 908
Bacteriology, Bergey manual of determinative, review of phytopathogenic forms in 5th ed., 128
Bacteriosis, walnut, control by insoluble copper sprays, 826
Bacterium (See also Phytomonas)
Bacterium angulatum, infection as influenced by water-soaking, 2, 889
dieffenbachiae, n. sp., leaf spot of Dieffenbachia picta due to, 961
malvacearum, on cotton seeds, as influenced by H_2SO_4 treatment, 879
medicaginis var. phaseolicola, seed treatment of beans for, 19
phaseoli, seed treatment of beans for, 19
solanacearum, tobacco wilt, control by rotation, 22
weeds as reservoirs of, 22
tabacum, infection as influenced by water-soaking of leaves, 889
tumefaciens, colchicine effects on tumors induced by, 221
BAIN, DOUGLAS C., 879
BAINES, R. C., 652, 654
BALDWIN, I. L., (852)
Barberry, eradication, results in Penna., 24
Bark beetles, blue-staining fungus associated, on pine, 556
Bark necrosis, internal apple-, as boron deficiency disease, 10
BARKER, H. D., (751), (752)
Barley, covered smut, chlorine-gas seed disinfection for, 913
as influenced by environment, 22
Fusarium spp. in diseases of, 7
loose smut, infection factors, 717
powdery mildew, control by water sprayed under pressure, 289
variety resistant to rust and other diseases, 209
BARRETT, J. T., 822
BARSS, HOWARD P., 381, (293)
Basisporium gallarum, corn arrested axillary shoots induced by, 656
Basswood, American, Nectria canker, growth and mortality relations, 23
BATTEN, E. T., 18
BAUR, KARL E., (436), (825)
Bean, broad, mild-mosaic virus (new) of, 448
mild-mosaic virus, host range, transmission, etc., 448
Sieva Lima, pea viruses infecting, 25
Windsor broad, pea viruses infecting, 25
Beans (See also Phaseolus)
anthracnose resistance, factorial interpretation, 1
bacterial blight, seed treatment for, 19
nodule hyperauxony, 15
phyllody in varieties, morphology and anatomy, 823
powdery mildew, control by water sprayed under pressure, 289
resistance inheritance, 824
strains and host range, 824

Rhizoctonia stem rot, host-parasite relations as influenced by nutrients, 1
rust, infection in relation to moisture, 933
physiologic races, 9, 820
web blight (Corticium microsclerotia), 559
BECKER, HANNAH (204)
Beech (See Fagus)
Beet (See also Sugar beet)
garden, internal black spot, etiology, 120
Begonia tuberhybrida, leaf-spot bacteriosis of, 633
BENNETT, C. W., 1, 422
Bentonite sulphur, fused, colloidality, 8
Benzol, toxicity of vapors, mechanism, 186
to tobacco seedlings vs. Peronospora tabacina, 177
vapor determination in air mixtures, 179
BERGEY, DAVID H., 128
BERGMAN, H. F., 545
Betula, butt and trunk rot due to Daedalia, 3
Nectria cankers, avenues of entry, 353
BEVER, WAYNE M., 863
Bioclimatics, new subsection of Biological Abstracts, 1004
Biographies and necrologies, George, Doroth Chauncey, 389
Meier, Fred Campbell, 293, 389
report, 380
Biological Abstracts, bioclimatics-biometeorology as new sub-section of, 1004
Biological control, of mistletoe, 995
of nematode injury to plants, 596
of root-rot fungi, 750
Biometeorology, new subsection of Biological Abstracts, 1004
Biometry, analysis of variance, review, 657
generalized standard errors in wheat-bunt tests, review, 657
statistical analysis of replicated experiments, 657
Birch (See Betula)
BLACK, L. M., 2, 321
Black locust (See Robinia pseudoacacia)
Black raspberry, mosaic, aphid transmission, 825
Black rot, kale, control, 19
Black spot, beet internal-, etiology, 120
rose Diplocarpon-, 315
Blackberry (See also Rubus)
stamen blight due to Hapalosphaeria deformans, 829
Blackfire, tobacco, as influenced by water-soaking of leaves, 884
Blackleg, potato, bacteria associated, 831
Black-raspberry, mosaic, aphid transmission, 825
BLAISDELL, DOROTHY J., 2, (24), 761
BLANK, LESTER M., 2
Blight(s), Antirrhinum Botrytis-, trichomes as infection courts for, 651
azalea and rhododendron bud and twig, Sporocybe azaleae causing, 517
bean bacterial, seed treatment for, 19
bean web-, Corticium microsclerotia causing, 559
blackberry stamen-, Hapalosphaeria deformans causing, 829
blueberry twig-, 136
camellia blossom-, Sclerotinia sp. causing, 824
cedar, in nurseries and its control, 14
celery, dissemination on clothing, 553
corn bacterial-, new host, 6, 284
invasiveness as influenced by nitrogen supply, 17
hemlock twig-, Melampsora farlowii causing, 463
of maple spp., phthiaceous fungus associated, 11
pea Septoria-, varietal reactions to, 25
peach Coryneum-, control by Bordeaux-petroleum oil emulsion, 828
pine brown-spot needle-, 1076
red cedar Phomopsis, in nursery stock, source and control, 991
tomato early-, control by cuprous oxides, 315
delayed spraying for, 11
wheat seedling-, seed treatments for, 16
Blind pocket, citrus, relationships, 6
Blister rust, white pine (See Cronartium ribicola)
BLODGETT, EARLE C., 651
Blossom blight, camellia, Sclerotinia sp. causing, 824
Blossom-end rot, tomato, environal- metabolic factors in development, 7
Blotch, potato tuber-, distribution, 932
Blueberries, Phomopsis twig blight, 136
powdery mildew on cultivated, varietal reactions to, 545
Blue-staining fungus, associated with Ips lecontei on Pinus ponderosa, 556
Blue-stem, potato, suberin and suberized deposits in, 278
Boehmeria drummondiana, Corticium microsclerotia on, 560
Boll rots, cotton-, fungi associated, 753
BONDE, REINER, 831
Book reviews (See under Reviews)
Bordeaux mixture, cucumbers injured by, 751
fungicidal value as influenced by petroleum-oil emulsion, 828
as influenced by petroleum-oil emulsion, 828
as peach summer fungicide, 11
in sour cherry leaf-spot control, 13
Boron deficiency, apple, 10
beet internal leaf spot in relation to, 120
prune drought spot in relation to, 650
Botrytis, Antirrhinum blight, trichomes as infection courts for, 651
Boysenberry, stamen blight due to Hapalosphaeria deformans, 829
BOZOVAISKY, LUBEN (242), (751)
Brachychaeta sphaeeolata, as Elsinoë solidaginis new host, 970
BRADFORD, F. C., (1002)
BRAUN, ARMIN C., 2
BREAKEY, EDWARD P., (825)
"Breaking," in tulip, first record (?), 1001

Breeding, beans, for anthracnose resistance, 1
cotton, for wilt resistance, 753
cowpeas, for disease resistance, 826
cucumber, for bacterial wilt resistance, 996
raspberry, for mosaic resistance, 647
snapdragon, for rust resistance, 18
sweetclover, for root-rot resistance, 909
tomato, for disease resistance, 215, 757
for Fusarium wilt resistance, 25
for leaf-spot resistance, 750
watermelon, for wilt resistance, 5
wheat, for leaf-rust resistance, 965
Brevity, in writing, 292
BRIERLEY, PHILIP, 3
BRINKEHOFF, LLOYD A., 823, (827)
Bromus inermis, Septoria leaf spot and susceptibility of inbred lines, 1, 554
Brooming, of black locust and other trees, virus induced, 8
Browallia speciosa, Solanum virus-4 in, 170
BROWN, H. G., 753
BROWN, NELLIE A., 221,
Brown rot, peach, control, 11
of stone fruits, apothecia destroyed by calcium cyanamid, 436, 825
distribution in California, 346
Sclerotinia spp. in California causing, 337
BRUYN, HELENA L. G. de, 3
Buckskin disease, cherry, control, 20
virus passage in relation to resistant stock, 20
Bud blight, azalea and rhododendron, Sporocybe azaleae causing, 517
Bulb rot, narcissus, aluminum sulphate control, 23
Bunt, chlorine gas seed disinfection for, 913
forage grass susceptibility, 575
wheat, chlorotic leaf mottling due to, 456
generalized standard errors in tests with, review, 657
reinoculation of resistant varieties with purified physiologic races, 863
BURKHOLDER, WALTER H., 128
BUTLER, KARL D., (624)
Butt rot, of hardwoods, Daedalia causing, 3
Buttercup, Persian, root rot due to Pythium spp. in, 828
B-virus, potato, on American vs. European varieties, 59

Cabbage maggot, as vector of Erwinia carotovora, 849
Cadophora, pines infected by spp. of, 1036
Calcium cyanamid, effect on population of Taeniothrips in orchards, 825
Sclerotinia apothecia destroyed by, 436, 825
CALDIS, P. D., 823
Calico, potato, 6,
differentiation and classification, 928
-Solanum virus 10, differentiation and classification, 928
California vine disease, transmissible disease resembling, 10
Calomel, potato-scab and scurf control by, 13
Camellia japonica, blossom blight due to Sclerotinia sp., 824
CAMPBELL, W. A., 3
Canada-streak, potato, 6,
differentiation and classification, 921
geographical distribution, 932
Cankers, apple perennial, as influenced by wound dressings, 739
basswood Nectria-, growth and mortality relations, 23
beech Polyporus-, sterile conks associated, 3,
cypress, as influenced by sea spray, 823
Nectria-, in hardwood trees, 351
peach, fungi involved, 10
pine blister rust-, development on young white pines, 1067
sugar beet dry rot-, 793
Canning peas, freezing injury to, 188
Caperonia castaneaefolia, leaf blotch (Ciboria associated), 749
Capsicum annuum, Solanum virus-4 in, 170
Verticillium wilt, 359
Carbon bisulphide, Armillaria mellea control in orchard soils by, 827
control of soil pathogens and weeds by, 25
nematode control in greenhouse by, 11
Carbon dioxide, evolution from soil, relation to black root rot of flue-cured tobacco, 751
Carex frankii, leaf blotch (Ciboria associated), 749
CAROSELLI, N., (11)
Carrot, phyllody in, 823
Carrot-rust fly, as vector of Erwinia carotovora, 849
CARTER, WALTER, 285
CASSELL, ROBERT C., 4, (22)
Castanea, Cronartium spp. spontaneous on, 998
North American spp. of Cronartium on, 998
Castanopsis, North American spp. of Cronartium on, 998
Cataphoresis, fungicide adherence properties determined by, 19
Cedar, blight, in nurseries and its control, 14
red (See Juniperus virginiana)
Celery, blights, dissemination on clothing, 553
pseudo-calico, host range, 824,
new virosis, 824
Septoria leaf spot of, pycnidial and spore numbers in, 646
virus diseases, 824
insect transmission, 824
viruses, properties, 824
CELINO, MARTIN S., (512)
Cephalosporium, elm die-back due to, 371
Ceratostomella, pine-staining spp., 1034, 1035, 1036
fimbriata, crop rotations in relation to, 750
ips, transmission by bark beetles, 556, 1049
pluriannulata, hardwoods stained by, 1037

ulmi, effect on elm spp., 23
eradication, scouting and sampling elms as aid to, 529
fructification on elms in England, 551
infection in relation to common elm borer, 547
in Scolytus galleries, 551
Cercospora, peanut leaf spots due to spp., control by dusts and sprays, 18, 751
strawberry leaf spot due to apparently new, 19
apii, dissemination on clothing, 553
vexans, strawberry Cercospora leaf spot differing from that due to, 19–20
zebrina, sweetclover seed parasitized by, 912
Cereals, Fusarium spp. in diseases of, 7
rusts of, ecology, 24
nature of resistance to, 672
seed-borne microorganisms, epidemiology of, 8
seed disinfectants compared, 16
smuts, ecology of, 22
weevils in, chlorine gas seed disinfection for, 913
Ceresan, sugarbeet damping off control by, 15
Charcoal rot, cowpea, breeding for resistance, 826
Chemotherapy, of plant diseases, 978
Cherry (See also Prunus)
buckskin disease, control, 20
virus passage in relation to resistant stock, 20
Coccomyces leaf spot, control by fungicides, 5, 13
Sclerotinia spp. on, in Wash., 825
sour, new bud-transmissible virus(?) disease, 821
Coccomyces leaf spot control by spraying, 13
sweet, gum-pocket formation in wood, microchemistry, 20
CHESTER, K. STARR, 4, 962
Chestnut (See Castanea)
Chinquapin (See Castanopsis)
Chlorine gas, as seed disinfectant, 913
Chloropicrin, control of soil pathogens and weeds by, 25
kale black-rot control by, 19
nematode control in greenhouse by, 11
Chlorosis, citrus zonate-, relationships, 6
sour cherry, virus(?)-induced, 821
Chlorotic leaf mottling, of wheat, bunt causing, 456
Choanephora cucurbitarium, cowpeas attacked by, 898
hosts, 898
CHUPP, CHARLES, (382)
Ciboria, leaf blotch of various hosts associated, 749
rice leaf blotch (new) associated with, 749
Cirsium nuttallii, Corticium microsclerotia on, 560
Citrus, infections variegation, 911
leprosis, lesions resembling on laurel oak, 907
relationships, 6
Phytophthora on, isolation method, 290
psorosis, registration of trees inspected for, 825
relation to other virus-like effects, 6
Cladosporium, hardwoods stained by, 1037
pines infected by, 1036
fulvum, tomato leaf mold due to, resistant red tomato, 9
CLARK, A., 657
CLARK, C. F., (362)
Claviceps paspali, on Paspalum spp. (including new hosts), 365
CLAYTON, C. N., (13), (821)
CLAYTON, E. E., 5
Clonal varieties, diseases in relation to, 9
Club root, as influenced by soil pH, 9
Clover, red (See Trifolium pratense)
Clovers, susceptibility to Pleospora lycopersici vs. Macrosporium sarcinaeforme, 541
Coccomyces hiemalis, control of cherry leaf spot due to, 5, 13
COCHRAN, L. C., (645)
COCHRAN, W. G., 657
Cod-liver oil, fungicidal value, 5
Colchicine, effect on bacterial vs. auxin-induced plant tumors, 221
Cold, resistance to, in oat varieties, as influenced by crown and stem rusts, 763
Collar rot, cotton internal, Thielaviopsis causing, 751
Colletotrichum falcatum, setae produced in culture, 7
lindemuthianum, resistance of beans to, factorial interpretation, 1
COLWELL, WILLIAM E., (651)
Concentric ring blotch, citrus, relationships, 6
Conifers, damping off of seedlings controlled by chemical treatments, 14
Sclerotium bataticola infection of seedlings, 14
COOK, HAROLD T., 5, (19)
COOLEY, J. S., 5
Copper, fungicides (See Fungicides, copper)
Corky bark, citrus, relationships, 6
Corn, arrested axillary shoots, Basisporium gallarum causing, 656
bacterial wilt, as influenced by nitrogen supply in sweet vars., 17, 1051
motile bacterium in relation to, 1062
new host, 6, 284
crazy top of, characterization and distribution, 817
dent, kernel discoloration in inbred and hybrid lines, 905
ear infection by Basisporium gallarum, 656
oil, fungicidal value, 5
Penicillium oxalicum infection, mechanism, 237
seedling injury by Penicillium spp. and factors influencing, 231
smut, growth types and factors influencing, 473
Cornus, Corticium galactinum infecting, 5

Corticium galactinum, apple-tree white root rot due to, 5
hosts, 5
microsclerotia, n. comb., bean web blight due to, 559
hosts, 560
vagum, cotton seedlings infected by, 373
life history, 373
sugar beet dry-rot canker due to, 793
Coryneum beijerinckii, control by Bordeaux-petroleum oil emulsion, 828
cardinale, spore germination and mycelial growth as influenced by sea spray, 823
Cotton, Acala, nematode infestation and control, 752
angular leaf spot, persistence and distribution factors under irrigation, 754
seed treatment by H_2SO_4 for, 879
boll rots, fungi associated, 753
breeding, for wilt resistance, 753
delinting of seed, effect on seed treatments, 754, 755
Fusarium wilt, as influenced by: acidity, and nitrogen source and level, 756
nitrogen, phosphorus and potassium nutrition, 756, 757
unbalanced fertilization, 756
inoculation technique, 755
soil contamination, experimental technique, 756
virulence of fungus strains, 1, 21, 757
water-culture infection method, 23
girdling of stems, effect on root rot, 753
internal collar rot (Thielaviopsis), 751
Rhizoctonia solani perfect stage from seedlings, 373
root-knot, in Calif., 752
control, 752
in Tenn., 751
root rot, green manure crops for, 752
incidence, rainfall relations, 752
seed, from diseased bolls, as influenced by H_2SO_4 treatment, 879
storage factors after mercury dust treatments, 754
seed treatment, effect on seedling survival, 755
with mercury dusts, as influenced by type and period of storage, 986
by organic mercury dust and H_2SO_4, 755
tests, 754
seedling diseases, fungi associated, 753
tests of seed, permeability of normal and treated, 753
Verticillium wilt, control, 753
wilt/nematode relations, 752
varietal reactions to, 755, 974
Cottonseed oil, fungicidal value, 5, 24
Cover crops, green-manure, for root rot, 752
Cowpea, Choanephora cucurbitarium on, 898
Corticium microsclerotia on, 561
powdery mildew, resistance inheritance, 824
strains and host range, 824
resistance to diseases, breeding for, 826
CRALLEY, E. M., 757
Crazy top, corn, characterization and distribution, 817
CRESS, MAX, (4)
Crinkle, potato, on American vs. European varieties, 46
Crinkle leaf, celery, aphid transmission, 824
Crinkly leaf, citrus, relationships, 6
Cronartium, North American spp. on Castanea, Castanopsis, and Lithocarpus, 998
species spontaneous on Castanea, 998
ribicola, canker development on young white pines by, 1067
on currant, varietal reactions, 26
immunity in staminate clone of Ribes alpinum, 981
inoculum source for greenhouse studies, 644
life history on young white pines, 1067
resistance of Viking currant, cytological study, 26
Ribes punctatum susceptibility, 643
Crops, rotation (See Rotation)
Crown gall, effect on young apple trees, 21
as influenced by colchicine, 221
pathogenicity basis, 16
Crown rot, of sugar beets, comparison to dry-rot canker, 796
Crown rust, oats, developmental factors, 659
effect on cold resistance, 763
urediospore viability, 21
variety reactions to, 763
Crucifers, club root, as influenced by soil pH, 9
Cucumber, bacterial wilt, resistant hybrids, 996
Bordeaux injury to, 751
downy mildew, control by copper fungicides, 751
infection in relation to moisture, 933
mosaic, cross-protection tests with virus strains, 903
virus as component of Easter lily mosaic virus, 3
pea viruses infecting, 25
powdery mildew, control by water sprayed under pressure, 289
Cucurbits, Pythium aphanidermatum causing root and fruit rots of, 624
Culture media, effect on fungus virulence, 757
fungus inoculum effect on colony growth in, 592
for fungus spore germination and growth, 828
isolation, for Actinomycetes, 1000
liquid, renewal method for fungi, 895
peptone broth, growth substances produced by crown gall and related bacteria in, 16
poured-plate method, efficacy for bacterial plant pathogens, 852
for sporulation of fungi, 821
Culture technique, for fungus identification and classification, 827
single spore isolator, 212

Culture tubes, label for, 761
Cultures, transfer hood for, 908
CUMMINS, GEORGE B., (654)
Cupressus macrocarpa, canker fungus of, spore germination and mycelial growth as influenced by sea spray, 823
Curly top, phyllody in plants with, 823
in tobacco, recovery from and acquired tolerance to, 743
virus, antibody-like substance in Turkish tobacco infected with, 828
-virus-1, concentration in root tips of resistant vs. susceptible sugar beet vars., 825
Currant, blister rust, resistance of Viking var., cytological study, 26
susceptibility of Andien sp., 643
varietal reactions, 26
mosaic, new virus disease, 369
mountain- (See Ribes alpinum)
Viking variety, leaf anatomy, 29
C-virus, potato, on American vs. European varieties, 61
Cyamopsis, Erysiphe polygoni on, 824
Cyclosis, citrus, relationships, 6
Cynodon dactylon, leaf blotch (Ciboria associated), 749
Cypress, canker fungus, spore germination and mycelial growth as influenced by sea spray, 823

Dactylaria candida, nematode injury restricted by, 600–,
thaumasia, nematode injury restricted by, 600–,
Dactylella ellipsospora, nematode injury restricted by, 599–,
Daedalea unicolor, butt and trunk rots of hardwoods due to, 3,
DAINES, ROBERT H., 5
Daldinia concentrica, hardwood decay by, 2
Damping off, in conifer seedlings, control by chemical treatments, 14
control by cuprous oxides, 306
in eggplants, tomatoes and peppers, seed and soil treatments for, 750
electrical pasteurization of soil for, 18
of mushrooms, Fusaria associated, and control, 728
pine distribution in relation to, 367
by Rhizoctonia solani, determination of physiological races, 609
sugar-beet, quasi-factorial vs. randomized-block designs of experiments on, 637
seed treatments for, and factors influencing, 15
DANA, B. F., 823
DARKIS, F. R., 103, (177)
Datura, differential host of tomato tip blight, 161
stramonium, Solanum virus-4 in, 169
DAVIDSON, ROSS W., (3), 5
DAVIS, WILLIAM C., (14), 991
DAVIS, W. H., 517
DECKER, PHARES, (204)
Deficiency diseases, beet internal black spot, 120
boron, in apple, 10
in prune, 650
grass degeneration (new), 95
pea marsh spot due to manganese deficiency, 3
Delphinium, aster yellows in, incubation period and leaf-hopper transmission, 826
DENNIS, R. W. G., 168
DIACHUN, STEPHEN, 231, (884)
Dieback, elm Cephalosporium-, 371
-virus of pea, 25
Dieffenbachia picta, bacterial leafspot of, 19, 956
DIMOCK, A. W., 823
Diplocarpon earliana, strawberry scoreh due to, new Mycosphaerella leaf spot resembling, 20
rosae, control by cuprous oxides, 315
Diplodia gossypina, incidence on cotton, 754
natalensis, hardwoods stained by, 1037
pines stained by, 1034
zeae, scolecospores produced in culture, 67
Diseases, clonal varieties in relation to, 9
control by toluene compounds, 978
deficiency (See Deficiency diseases)
dissemination, on clothing, 553
forest nursery-, 14
forest tree-, 915
of fruits, book review, 915
of hops, book review, 915
of ornamentals, book review, 1002
plant, handbook, review, 558
new etiological terminology, 459
resistance to (See under Resistance)
seed-borne of cereals, epidemiology, 8
chlorine-gas disinfection for, 913
sugar cane, book review, 220
virus (See also under Virus)
annotated list (China), 459
first record (?) of plant-, 1001
virus and virus-like, of trees, 8
Disinfectants, seed (See also Fungicides; Seed treatment; etc.)
seed-, chlorine gas as, 913
Dissemination, disease (See under Diseases; Insects as vectors; and specific entries)
DOAK, K. D., (14)
Dogwood, Corticium galactinum infecting, 5

Doidella teres, Corticium microsclerotia on, 560
Dolichos, Erysiphe polygoni on, 824
DOOLITTLE, S. P., (757), 996
Dothiorella ulmi, elm die-back due to, 371
Downy mildew (See under Mildew)
DRECHSLER, CHARLES, 391, 1005
Drought spot, of prunes, as boron deficiency, 650
Dry-rot canker, sugar beet-, 793
DUGGAR, B. M., (16)
Duke Forest, fungi of, book review, 915
DUNDAS, B., 820, 824
DUNEGAN, JOHN C., 72
Dutch elm disease, effects on elm spp., 23
eradication, scouting and sampling elms as aid to, 529

fungus fructifications in England, 551
infection in relation to common elm borer, 547
DYKSTRA, T. P., 6, 40, 917

Early blight, tomato, control by cuprous oxides, 315
delayed spraying for, 11
Easter lily, mosaic complex of, two viruses in, 3
Echinochloa colona, Corticium microsclerotia on, 560
crusgalli, corn crazy-top-like symptoms in, 820
leaf blotch (Ciboria associated), 749
Ecology, of cereal stem rusts, 24
of fungi staining wood, 1046
of fungus growth types, 473
of oats loose smut, 22
of red pine mycorhizas and parasites, 24
of smut diseases of cereals, 22
EDGERTON, C. W., (19)
EDSON, H. A., 377, 378
Eelworms (See nematodes, and under scientific names)
EGGERS, VIRGINIA, (15)
Eggplants, damping off, seed and soil treatments for, 750
EHRLICH, JOHN, 459, 908
Electrical apparatus, for soil pasteurization, 18
Electrodialysis, fungicide adherence properties determined by, 19
Elements, minor, deficiency disease of grass in relation to, 95
Eleusine indica, Corticium microsclerotia on, 560
ELLIOTT, CHARLOTTE, 6, 284
ELLIS, DON E., 556, 995
Elm (See Ulmus)
Elm borer, common, as vector of Dutch elm disease, 547
Elsinoë solidaginis, new hosts and distribution, 970
Elymus, Ustilagos on spp. of, 490
Emelista tora, Corticium microsclerotia on, 560
Endoconidiophora, pines infected by spp., 1036
coerulescens, hardwoods stained by, 1037
moniliformis, hardwoods stained by, 1037
Entomophthora, differentiation method, 9
strain from apple leaf hopper, 9
Ergot, of Paspalum spp., 365
Errata, for Phytopathology (Vol. 28), 390
Errors, generalized standard, in wheat-bunt tests, review, 657
Erwinia. (See also Bacillus.)
amylovora. (See also specific hosts.)
exudate from infections, nature and properties, 142
survival in dried vs. moist ooze, 142
carotovora, insect dissemination, 849
potato blackleg, seed-piece decay and soft rot due to, 831
strains compared, 831
Erysiphe cichoracearum, control by water sprayed under pressure, 289
graminis, control by water sprayed under pressure, 289
hordei, barley variety resistant to, 211
polygoni, control by water sprayed under pressure, 289
culture trials and factors influencing, 828
resistance in beans and cowpeas, inheritance, 824
strains and host range, 824
Etiology, new terminology, 459
Euonymus, powdery mildew, control by water sprayed under pressure, 289
Eutettix tenellus, virus activity inhibited by juices of, 324
Experimental technique (See under Technique)
EZEKIEL, WALTER N., 752, 753

Fagus, cankers and associated sterile conks of Polyporus on, 3
FAWCETT, H. S., 6, (290), 907, 911
FELIX, E. L., 6
Ferrous sulphate, in conifer damping off control, 14
Fertilization, of corn, effect on bacterial wilt, 1051
of cotton, effect on Fusarium wilt, 756, 757
of Para rubber tree, effect on Fomes lignosus infection, 14
of red cedar, effect on blight incidence in nursery stock, 992
of sugar beet, effect on Sclerotium infection, 15
Ficus carica, Corticium microsclerotia on, 560
Field experiments, quasi-factorial vs. randomized-block designs for, 637
Fig, Corticium microsclerotia on, 560
Fiji disease, sugarcane POJ 2878 behavior to, 512
virus, transmission by Perkinsiella vastatrix, 512
Filtration, of tobacco-necrosis plant juices, 83
Fire blight, bacterial survival in dried vs. moist ooze, 142
exudate, nature and properties, 142
Firmiana platanifolia, Corticium microsclerotia on, 560
FISCHER, GEORGE W., 490, 575
Flowers, garden and greenhouse, diseases and pests of, book review, 1002
Flue-curing, of tobacco, effect on infectivity of mosaic virus-1, 242
temperatures for, 246
FOLSOM, DONALD, 6
Fomes lignosus, on Hevea, 14
as influenced by fertilizers, 14
physiologic races, pathogenicity compared, 14
Forage grasses, smut susceptibility, 490
FORBES, IRVIN L., 7, 659
Forestry, clonal varieties for tree planting, parasites in relation to, 9
nurseries, disease developments in, 14
pathology, book review, 915

Formaldehyde, effect on black locust emergence, 14
Fortunella margareta, water damage to fruit, 214
FOSTER, A. C., 7
FOSTER, H. H., (5)
FRAMPTON, VERNON L., 7, 495
Frankliniella schultzei, as vector of pineapple yellow-spot virus, 286
Fraxinus americana, virus-induced blooming of, 8
nigra, virus-like symptoms on, 8
FREAR, DONALD E. H., (993)
Freezing, injury to canning peas by, 188
FREITAG, JULIUS H., 824
Frenching, tobacco, thallium injury in relation to, 502
Frosts, canning-pea injury by late spring-, 188
Fruits, clonal varieties in relation to diseases, 9
diseases, book review, 915
production fundamentals, book review, 1002
rots, Pythium aphanidermatum causing, 624
Fungi, autolysis in crosses of Sphaerotheca, 14
of Duke Forest, book review, 915
growth substances in relation to, 592
growth types as influenced by genetic vs. environal factors, 473
of India, book review, 374
inheritance in, 473
inoculation technique, water-culture method, 23
inoculum effect on colony growth, 592
life histories, unusual features in Phymatotrichum, 498
nematode injury to plants restricted by, 596
nutrition, 592
physiologic races (See Physiologic races)
pythiaceous, in maple blight, 11
renewed liquid cultures of, 895
root-rot, biological control, 750
soil treatments for, 750
soil contamination technique, 755
spore germination trials, factors influencing, 828
of fungicides, variation factors analyzed, 16
spore isolator for, 212
sporulation, of, culture media for, 821
transfer hood for cultures, 908
virulence, as influenced by culture media, 757
wood-destroying, 1036
wood-staining, relative importance and seasonal prevalence in United States, 1031
Fungicide(s), adherence properties, determination by cataphoresis, 19
for bean bacterial blight seed treatment, 19
bentonite sulphur, colloidality, 8
benzol vapors, for tobacco downy mildew, 186
Bordeaux, cucumber injury by, 751
as influenced by petroleum-oil emulsion, 828
for peach in summer, 11
calcium cyanamid, for brown rot apothecial destruction, 436, 825
calomel, for potato scab and scurf, 13
carbon bisulphide, for Armillaria in soils, 827
for cereal seed treatment, comparisons, 16
for cherry leaf spot, 5, 13
chlorine-gas, as seed disinfectant, 913
chloropicrin, for kale black rot seed treatment, 19
for soil pathogens, 25
cod-liver oil as, 5
for conifer damping off, 14
copper, adherence, factors in, 19
properties as determined by chemical analysis and cataphoresis, 19
compatibility with arsenical and nicotine-oil sprays, 12
cucumber downy-mildew control by, 751
insoluble, for walnut bacteriosis, 826
oils as supplements to, 24
for peanut leaf spots, 751
red oxide of, for sugar-beet damping off, 15
for cotton seed treatments, 754
effect on seedling survival, 755
cottonseed oil as, 24
cuprous oxides, particle size and color in relation to protective value, 303
yellow, small particle size, 9
dusts, measuring interfacial friction of seeds treated by, 200
formaldehyde, effect on black locust emergence, 14
heavy-metal, effects on Phymatotrichum, 2
lime-sulphur causing apple yield reductions, 6
liquid lime-sulphur, standardized procedures important in diluting, 993
mercuric chloride-alcohol-acetic acid, for bean bacterial blight seed treatment, 19
mercuric oxide, for narcissus bulb rot, 23
mercury dusts, cotton-seed treatments as influenced by type and period of storage, 986
organic, cotton seed storage factors after use, 754
oils (animal and vegetable) as, 5, 24
paradichlorobenzol, as tobacco downy-mildew eradicant, 216
for peanut leaf spot, copper and sulphur, 18, 751
phenothiazine and derivatives as, 7, 700
for potato seed treatment, Rhizoctonia tuber penetration in relation to, 759
seed-disinfectant, for cereals, 16
chlorine gas as, 913
soil-, 25
for narcissus bulb rot, 23
for root-rot fungi, 750

for sour cherry leaf-spot control, 13
spore-germination tests, variation factors analyzed, 16
spray injury by, as influenced by oil supplements, 24
spray residue removal, 12
spray supplements, oils as, 24
spraying by airplane, in peach orchards, 823
for sugar-beet seed treatment, 15
sulphur, elementary, particle size, 8
fused bentonite-, colloidality, 8
for peanut leaf spots, 751
for sweet-potato soil rot, 750
sulphuric acid, in cotton-seed treatment, 879
toluene compounds as, 978
for tomato early blight, 11
vegetable oils as, 5
volatile, toxicity test method, 641
wood preservatives, acceleration of toximetric tests by soil, 901
wound dressings, effect on apple perennial canker, 739
Fusarium, antagonism between Agaricus and, 728
Arceuthobium disease due to (?), 995
identification and classification, culture methods in relation to, 827
incidence of spp. on cotton, 754
mushroom damping-off due to, 728
narcissus bulb rot due to, mercuric oxide soil treatment for, 23
pines infected by, 1036
species associated with cereal diseases, 7
avenaceum, cereal disease association, 7
suberin and suberized deposits in potatoes infected with, 278
batatatis, crop rotations in relation to, 750
bulbigenum var. lycopersici, host resistance and pathogenicity, study technique, 945
resistant tomato hybrid, 757
culmorum, cereal-disease association, 7
equiseti, cereal-disease association, 7
lycopersici, chemical soil treatment against, 25
cultural characters and pathogenicity of strains, 24
resistance to, time-weighting method of comparing, 873
tomato varietal resistance to, 24, 25, 871
moniliforme, incidence in cotton disease, 754
niveum, watermelon variety resistant to, 5
oxysporum var. aurantiacum, cereal-disease association, 7
poae, cereal-disease association, 8
scirpi var. acuminatum, cereal-disease association, 8
vasinfectum, on cotton, as influenced by: acidity and nitrogen source and level, 756
nitrogen, phosphorus and potassium nutrition, 756, 757
unbalanced fertilization, 756
inoculation technique, 755
nematodes in relation to, 752
varietal susceptibility to root-knot nematode plus, 974
soil contamination, experimental technique for, 755
variability, 21, 755
virulence, differences in isolates, 1
as influenced by culture media, 757
strains compared, 757
variability in, 21
tracheiphilum, cowpea breeding for resistance to, 826
water-culture infection method for, 23

Garden, -flowers, diseases and pests, book review, 1002
GARDNER, V. R., 1002
GARREN, K. N., (915)
GEMMELL, ALAN R., 95, 287
GEORGE, DOROTH CHAUNCY, biographical note on, 389
Germination tests, of spores vs. fungicides, variation factors analyzed, 16
Gibberella saubinetii, barley variety resistant to, 210
GIDDINGS, N. J., 649
GINGRICH, N. S., (17)
Girdling, of cotton stems, effect on root rot, 753
Gleditsia, virus induced brooming in, 8
Gleosporium perennans, apple canker due to, as influenced by wound dressings, 739
Glomerella cingulata, fungicidal action of phenothiazine and derivatives on, 701
gossypii, incidence in cotton boll and seedling diseases, 753
Gloxinia, Phytophthora cryptogea infecting corms, stems and leaves of, 17
Glycine apios, Corticium microsclerotia on, 560
GODFREY, R. K., (22)
Goldenrod (See Solidago)
GOLDSWORTHY, M. C., 7, 700, 1002
GOODWIN, M. W., (12)
GORDON, W. L., 7
GOTTLIEB, MANFRED, 624
GRAINGER, JOHN, 441
GRANT, THEODORE J., 8, 351
Granville tobacco wilt (See Bacterium solanacearum)
Grapes, Plasmopara viticola overwintering in wild, 822
red-leaf disease, control by mite control, 10
transmissible disease resembling so-called Calif. vine disease, 10
Graphium, pines infected by spp. of, 1036
rigidum, hardwoods stained by, 1037
Grasses, corn crazy-top-like symptoms in, 820
"degeneration" a new deficiency disease of Metropolitan bent, 95
ergot of Paspalum spp., 365
leaf blotch (Ciboria associated), 749
Puccinia rubigo-vera tritici inoculations to spp. of, 965
root knot incidence and control on, 752
Septoria leaf spot of Bromus inermis, 1, 554

smut susceptibility of forage-, 490, 575
stem rusts on cereals as influenced by incidence of, 24
GREANEY, F. J., 8, (16)
GREEN, E. L., (7), (700)
Green manures, for root rot, 752
Greenhouse, experiments, quasi-factorial vs. randomized-block designs for, 637
flowers, diseases and pests, book review, 1002
GROSS, P. M., (103), (177)
GROVES, A. B., 8
Growth substances, colchicine effects on plant tumors induced by, 229
crown-gall and related bacteria producing on peptone broth, 16
fungus relataions to, 592
in root nodules of Phaseolus, increase in, 15
GUBA, E. F., 9
Gum disease, citrus concave-, relationships, 6
Gum pockets, in sweet cherry wood, microchemistry of, 20
Gymnosporangium, apple rusts due to spp., varietal susceptibility relations, 18
clavipes, comparisons with other apple rusts, 801
globosum, comparisons with other apple rusts, 801
juniperi-virginianae, comparisons with other apple rusts, 801
on red cedar, infection time and control as influenced by aeciospore germinability and dissemination, 812

HAASIS, FRANK A., (23), 890
Hadromycosis, pepper, Verticillium causing, 359
HAENSELER, C. M., 9
HAHN, GLENN GARDNER, 643, 981
Hairy root, effect on young apple trees, 21
Hairy sprout, potato, non-transmissible form, 21
HANSEN, H. N., 824, (827)
Hapalosphaeria deformans, blackberry stamen blight due to, 829
Hardwoods, butt and trunk rot due to Daedalia, 3
decay by Ustulina and other Ascomycetes, 2
fungi infecting, 1037
Nectaria cankers on, avenues of entry, 351
HARRAR, J. G., 9
HARRISON, GEORGE J., 755, (752), (753)
HART, HELEN, 978
HARTER, L. L., 9
HARTLEY, CARL, 9, 657
HASKELL, ROYAL J., 293, (382)
Hawthorn, rust of, comparison with other rusts affecting apple, 801
HEDGCOCK, GEORGE G., 998
Helianthus annuus, rust, infection in relation to moisture, 933
Helminthosporium, pines infected by spp., 1036
geniculatum, hardwoods stained by, 1037
gramineum, barley variety resistant to, 211
Hemlock, rust due to Melampsora farlowii and control, 463
poison-, ringspot aphid transmitted, 824
HEPTING, GEORGE H., 463
HERRICK, J. ARTHUR, 504
Heterodera marioni (See also Nematodes)
chemical soil treatment for, 25
on cotton, 751
varietal susceptibility to Fusarium wilt plus, 974
cowpea breeding for resistance to, 826
effect on tomato resistance to Fusarium wilt, 871
host specialization, factors in, 752
pineapple injury restricted by fungi, 596
on tobacco, as influenced by transplanting dates, 21
on tomatoes, greenhouse control, 11
in Tenn., 751
radicicola, crop rotations in relation to, 750
schachtii, potato-root infestation by, 287
HEUBERGER, JOHN W., (11), 303
Hevea brasiliensis, Fomes lignosus on, 14
HEWITT, J. LEE, 825
HEWITT, WM. B., 337, (373)
Hicoria, Nectria cankers of, avenues of entry, 353
Hickory (See Hicoria)
HILDEBRAND, E. M., 10, 142, 369
HILL, H. H., (970)
HILL, L. M., 274
HIRT, RAY R., 1067
Hollow heart, potato, suberin and suberized deposits in, 280
HOLMES, FRANCIS O., 215, 431
Honeydew melons, Pythium root and fruit rot, 624
Honey locust, virus-induced brooming, 8
HOOKER, H. D., (1002)
HOPPERSTEAD, S. L., (12)
Hops, diseases, book review, 915
downy mildew, infection in relation to moisture, 933
Hordeum nodosum, Tilletia spp. infecting, 576
Ustilago spp. on, 490
HORSFALL, J. G., 9, 11, (200), (303)
Horticulture, fundamentals of fruit production, book review, 1002
Host-parasite relations, of Cronartium ribicola, 26
as influenced by nitrogen supply, 1051
in Rhizoctonia stem rot of bean, as influenced by nutrient variations, 1
Hot-water treatment, for peach phony disease, 12
HOUSTON, B. R., (15)
HOWARD, F. L., 11
HUBER, GLENN A., 436, (647), 825
HUMPHREY, H. B., 379, 385
HIRT, R. H., 11, 12
HUTCHINS, LEE M., 12
Hybrids, fungus-, autolysis in, 14
Hydrogen ion concentration, effect on: club root development, 9

cotton wilt, 756
plant pathogens, 750
Hyoscyamus niger, Solanum virus-4 in, 170
Hyperauxony, in bean nodules, 15
Hypoxylon, hardwood decay by sp. of, 2

Immunity, mountain currant clone, from blister rust, 981
in plants, material for study of, 749
potato, from X-virus, 362
sugar cane from viruses, virus classification tested by, 7
Immunization, from virus diseases in plants, 828
India, fungi of, book review, 374
Indoleacetic acid, colchicine effects on plant tumors induced by, 229
Infection courts, trichomes as, 651
Infections, bacterial, as influenced by water-soaking, 2
Inheritance, beans, to anthracnose, 1
cowpea, to powdery mildew, 824
fungi, 473
immunity from X-virus, in potato, 362
Inoculation chamber, for insect vector tests, 649
Inoculation technique, for fungus soil contamination, 755
for Fusarium, 755
water-culture method, 23
Inoculum, fungus-, effect on colony growth, 592
Insect pests, of ornamentals, book review, 1002
Insecticid(s), arsenical, compatibility with copper fungicides, 12
nicotine-oil, compatibility with copper fungicides, 12
removal from apples, 12
seed-disinfectant, chlorine gas as, 913
Selocide, in grape-mite and red-leaf control, 10
Insects, populations of individual plants, determination method, 20
virus activity inhibited by juices of, 321
Insects as vectors, aster yellows in delphiniums, by leaf hoppers, 826
black-raspberry mosaic, by aphids, 825
celery virus diseases, by aphids, 824
Ceratostomella ips, by bark beetles, 556, 1049
Dutch elm disease, by bark beetles, 551
by common elm borer, 547
Erwinia carotovora, 849
Fiji disease of sugarcane, by Perkinsiella castatrix, 512
inoculation chamber for, 649
mild mosaic of broad bean, by aphids, 450
pineapple yellow-spot virus, 286
potato purple-top, by aster leaf hopper, 14
raspberry mosaic, by Amphorophora rubi, 647
Sclerotinia fructicola, by thrips, 825
tomato tip-blight virus, by thrips, 159
viruses, as influenced by insect juices, 2
Internal bark necrosis, apple-, as boron-deficiency disease, 10
Internal black spot, garden beet-, etiology, 120
Ips, as vector of Ceratostomella ips, 556, 1049
Irrigation, cotton angular leaf spot spread by, 754
Isolation technique, for Actinomycetes, 1000
for Phytophthora spp., 290
single-cell, efficiency for bacteria, 852
single-spore isolator, 212
IVANOFF, S. S., 12, (616)

JACOB, H. E., (10)
JAMISON, CLYDE, (962)
Jasminum, rust on spp. of, 783
grandiflorum, rust on, 783
JENKINS, A. E., 970
JOHANN, HELEN, 67
JOHNSON, A. G., 380, 385
JOHNSON, E. M., (884)
JOHNSON, H. W., (15)
JOHNSON, JAMES, (2)
JOHNSTON, C. O., 456
JONES, FRED REUEL, 909, 912
JONES, L. R., 386
Juglans nigra, virus induced brooming, 8
regia, bacteriosis, control by insoluble copper sprays, 826
Juniperus virginiana, blight in nursery and its control, 14
Gymnosporangium juniperi-virginianae on, infection time and control as influenced by aeciospore germinability and dissemination, 812
Phomopsis blight in nursery stock, source and control, 991

KADOW, K. J., 12
Kale, black rot, seed treatment by chloropicrin for, 19
KEITT, G. W., 13, 821
KENKNIGHT, GLENN, 13, 1000
Kernel discoloration, in inbred and hybrid dent corn, 905
KERNKAMP, MILTON F., 473
KIENHOLZ, J. R., 13, 260
KING, C. J., 751, 752, 754
KING, THOMAS H., 14
KIRBY, R. S., (24), 376, 379, (382)
KLOTZ, L. J., 214, 290, (911)
KOEHLER, BENJAMIN, 817
KREUTZER, W. A., 629
Kumquat, water damage to fruit, 214
KUNKEL, L. O., 684

LACKEY, C. F., 825
LAMBERT, EDMUND B., 212
LAMIMAN, J. F., (10)
LASKARIS, THOMAS, 14
Latent mosaic, potato, immunity inheritance from, 362
LATHAM, DENNIS H., 14, (991)
Lathyrus odoratus, mild mosaic of broad bean infecting, 452
LAVIN, GEORGE I., (502)
LAWYER, LEWIS O., (827)
LEACH, J. G., 14, 204
LEACH, L. D., 15, (337), 373
Leaf blotch, rice, etc., Ciboria associated, 749

Leaf curl, virus induced, in China, 461
Leaf hopper, apple-, Entomophthora sp. from, 9
aster-, as vector of potato purple-top, 14
clover-, effect on virus of potato yellow dwarf, 2
Leaf hoppers, as vectors of: aster-yellows in delphiniums, 826
potato purple-top, 14
virus activity inhibited by juices of, 324
Leaf mold, tomato Cladosporium-, resistant red variety, 9
Leaf mottling, wheat chlorotic-, bunt causing, 456
Leaf roll, virus-induced, in China, 460, 461
Leaf spot(s), begonia bacterial, 633
black locust, 14
Bromus inermis, Septoria bromigena causing, 1, 554
celery, Septoria pycnidial numbers in, 646
cherry Coccomyces-, control, 5, 13
cotton angular, persistence and distribution factors under irrigation, 754
Dieffenbachia bacterial, 19, 956
fungi causing, media for sporulation of, 821
maple bacterial, 968
peanut Cercospora-, control, 18, 751
sour cherry Coccomyces-, control by spraying, 13
strawberry, Cercospora, 19
Mycosphaerella (new), 20
tobacco angular, infection as influenced by water-soaking, 2, 889
tomato, varietal reactions to Macrosporium and Septoria, 750
Leaves, tobacco, injury by water-soaking, 884
Lecanosticta acicola, perfect stage taxonomy, 1076
LECLERG, E. L., 267, 609, 637, 793
LEE, ATHERTON, 220
LEFEBVRE, C. L., 15, 365, (456), 898
Legume nodule bacteria, auxin increase in nodules induced by, 15
Legumes, mild mosaic of broad bean infecting other, 452
susceptibility to Macrosporium sarcinaeforme vs. Pleospora lycopersici, 541
LEHMAN, S. G., 755
Lemon, infectious variegation, 911
LEONARD, W. H., (657)
LEONIAN, LEON H., 592
Leprosis, citrus, lesions resembling on laurel oak, 907
relationships, 6
Leptographium, hardwoods infected by, 1037
pines infected by, 1036
Leptosphaeria pratensis, sweetclover seed parasitized by, 912
Lespedeza, bacterial wilt of spp., 15
LEUKEL, R. W., 913
LEUTRITZ, JOHN, Jr., 901
Lilium longiflorum, mosaic complex of, two viruses from, 3
LILLY, VIRGIL GREENE, (592)
Lime sulphur, apple yield reduction by spraying with, 6
liquid, commercial vs. homemade, 995
standardized procedures important in diluting, 993
LIN, K. H., 646
Linden, American, Nectria canker, growth and mortality relations, 23
LINDSAY, M. A., (752)
LINFORD, M. B., 596
LINK, GEORGE K. K., 15
LINN, M. B., 553
Linseed oil, fungicidal value, 5
Lippia lanceolata, anthracnose due to Sphaceloma lippiae, 654
Lithocarpus, North American spp. of Cronartium on, 998
LOCKE, S. B., 16
Locust, black (See Robinia pseudoacacia)
Lolium, Tilletia spp. infecting spp. of, 576
Lumber, fungi destroying, 1036
wood-staining fungi on, 1031
Lychnis alba, Corticium galactinum infecting, 5
viscaria, stem rot due to Phytophthora cactorum, 652
Lycium barbarum, Solanum virus-4 in, 171
Lycopersicum, Solanum virus-4 in spp. of, 170
chilense, as disease resistance source, 215
hirsutum, hybrid with L. esculentum mosaic-tolerant and wilt-resistant, 757
Lygus pratensis, potato wilt induced by, non-identity with purple-top, 14
Lysis, in Sphaerotheca sorghi crosses, 14

MCCALLAN, S. E. A., 16, 383
MCCORMICK, FLORENCE A., 371
MCCULLOCH, LUCIA, 956
MCFADDEN, E. S., 644
MCKINNEY, H. H., 16, (250)
MACLACHLAN, J. D., (1)
MCLEAN, RUTH, (103), (177)
MCNEW, GEORGE L., 17, 1051
MCREYNOLDS, D. K., 17
MCWHORTER, FRANK P., 651, 826

MACHACEK, J. E., (8), 16
MACKIE, W. W., 826
Macrosiphum, virus activity inhibited by juices of spp., 324
pisi, as vector of mild mosaic of broad bean, 451
Macrosporium, tomato leaf spot due to, varietal reactions, 750
sarcinaeforme, inoculations on tomato and legumes, 541
relationship to Pleospora lycopersici, 541
Macrosteles divisus, potato purple-top transmitted by, 14
virus activity inhibited by juices of, 324
Maggots, as vectors of Erwinia carotovora, 849
Maize (See corn)
Manganese, pea marsh spot due to deficiency, 3

Manures, green, for root rot, 752
Maple (See Acer)
Marsh spot, of pea, manganese deficiency causing, 3
MARTIN, J. P., 220
MARTIN, W. J., 17
MEIER, FRED CAMPBELL, biographical note on, 389
biography and list of publications, 293
portraits, 302 and facing p. 293
Melampsora farlowii, hemlock rust due to, 463
Melilotus, fungus parasites infecting seed of spp., 912
alba, mild mosaic of broad bean infecting, 452
Phytophthora root rot, breeding and selection for resistance, 909
susceptibility to Macrosporium sarcinaeforme vs. Pleospora lycopersici, 541
Melons, honeydew, Pythium root and fruit rot, 624
Melothria crassifolia, Corticium microsclerotia on, 560
Mercuric chloride,-alcohol-acetic acid, for bean bacterial blight seed treatment, 19
Mercuric oxide, narcissus bulb rot control by, 23
Mercury dusts, cotton-seed treatment by, as influenced by type and period of storage, 986
Mercury fungicides, organic, cotton-seed storage factors after use, 754
Metals, heavy, effect on Phymatotrichum omnivorum, 2
Meteorology, in new subsection of Biological Abstracts, 1004
Microsphaeria, blueberry powdery mildew due to sp. of, varietal reactions to, 545
MIDDLETON, JOHN T., 17, 541, (828)
MILBRATH, J. A., 156
Mildew, barley powdery-, control by water sprayed under pressure, 289
variety resistant to, 211
bean powdery-, resistance to, 824
strains and host range, 824
blueberry powdery-, varietal reactions to, 545
cowpea powdery-, resistance to, 824
strains and host range, 824
cucumber downy-, control by copper fungicides, 751
grape downy-, overwintering in wild grape, 822
hops downy-, infection in relation to moisture, 933
onion downy-, control, 18
culture trials of fungus, 828
infection in relation to moisture, 933
peach and rose forms of Sphaerotheca pannosa compared, 282
tobacco downy-, benzol vapor toxicity to, 177
control by: benzol vapors, concentration, 103
oil-copper combinations, 5
endemism, inoculum sources, dissemination and weather relations, 194
infection studies in flue-cured tobacco, 79
paradichlorobenzene as eradicant fungicide for, 216
in South America, 291
Mildews, downy, infection in relation to moisture, 933
oil sprays for, 24
powdery, control by water sprayed under pressure, 288
oil sprays for, 24
MILES, L. E., 754, 755, 974, 986
MILLER, J. K., (915)
MILLER, LAWRENCE I., (9), 18
MILLER, PAUL L., 18
MILLER, PAUL R., 753, 801, 812
MILLER, P. W., 826
Minor elements (See also Deficiency diseases)
new deficiency disease of grass in relation to
Mistletoe, biological control by fungus infection, 995
Mites, on grapes, control by Selocide, 10
rôle in red-leaf disease, 10
Mollusks, on ornamentals, book review, 1002
Monilochaetes infuscans, crop rotations in relation to, 750
Mosaic(s), annotated list (China), 459
black-raspberry, aphid transmission, 825
broad bean mild-, host range, transmission, etc., 448
new virus, 448
cucumber, cross-protection tests with virus strains, 903
currant, new virus disease, 369
Easter lily, two viruses in, 3
narcissus, white streak compared, 826, 890
potato, crinkle-, on American vs. European varieties, 46
latent-, immunity inheritance from, 362
leaf rolling, on American vs. European varieties, 49
mild-, on American vs. European varieties, 45
yellow group, compared and classified, 917
symptoms and virus properties, 6
raspberry, breeding for resistance to vector and, and factors in, 647
tobacco, virus, concentration as influenced by nitrogen nutrition, 22
detection in plant by rapid reagent-indicator method, 20
as influenced by insect juices, 2
inhibition by insect juices, 321
movement in tissues and factors in, 1
movement in tomato plants, 684
molecular weight of protein of, 495
physico-chemical study of protein of, 7
temperature relations of host and, 441
thermal inactivation, 20
virus-1, infectivity as influenced by flue-curing, 242

virus-6, purification and properties of protein of, 250
X-ray diffraction and molecular weight of virus protein prepared by Na_2SO_4 method, 17
tomato, hybrid tolerant to, 757
tulip, first record (?), 1001
turnip, virus inhibition by insect juices, 323
viruses, sugarcane immunity reactions in testing classifications of, 7
Mottle, potato, on American vs. European varieties, 57
Mottling, chlorotic leaf-, in wheat, bunt causing, 456
Mountain currant (See Ribes alpinum)
MUNCIE, J. H., (1000)
MUNDKÜR, B. B., 374
MURPHY, H. C., 763
MURRAY, C. W., (739)
Mushrooms, "damping-off," cause and control, 728
Muskmelons, quil, Pythium root and fruit rots of, 624
Mycology, review of "Fungi of India," 374
Mycorhizas (See also Pseudomycorhizas)
on pines, 14
on red pine, environal relations, 24
Mycosphaerella, strawberry leaf spot (new) due to sp., 20
fragariae, strawberry Cercospora leaf spot resembling that due to, 19
lethalis, sweetclover seed parasitized by, 912
platanifolia, sporulation induced by host-juice media, 821
Myzus persicae, virus activity inhibited by juices of, 324

Narcissus, Fusarium rot of bulbs, mercuric oxide as soil antiseptic for, 23
white streak, new virus disease, 826, 890
Nasturtium, differential host of tomato tip blight, 162
NEAL, D. C., 754, 755
Necrosis, apple, internal bark-, as boron deficiency disease, 10
potato net-, suberin and suberized deposits in, 280
potato top-, on American vs. European varieties, 61
tobacco, characteristic macromolecular substances in, 86
thermal inactivation of virus of, 20
ultrafiltration of juices of plants with, 83
virus, inhibition by insect juices, 323
properties, 84
Nectria, basswood canker due to, growth and mortality relations, 23
cankers of hardwoods due to, avenues of entry, 351
Needle blight, pine brown spot-, 1076
NELSON, O. A., (913)
NELSON, RAY, 18
Nematicides, for root knot and meadow nematodes, 752
Nematodes, in black locust nurseries, control by heat, 4
chemical soil treatment for, 25
on cotton, control, 752
infestation by wilt plus, varietal reactions to, 755
in Tenn., 751
varietal susceptibility to Fusarium wilt plus, 974
wilt in relation to, 752
cowpea breeding for resistance to, 826
electrical pasteurization of soil for, 18
estimation method for infestations, 287
meadow, on cotton and other plants, 752
on ornamentals, book review, 1002
pine infestations, 1036
pineapple injury restricted by fungi, 596
plant injuries by, measurement, 602
potato root infestation by, 287
root-knot, hosts and control in California, 752
soil populations, 601
stem-, oats and peas infested by, 827
on tobacco, as influenced by transplanting dates, 21
on tomato, effect on resistance to Fusarium wilt, 871
greenhouse control, 11
in Tenn., 751
Net necrosis, potato, suberin and suberized deposits in, 280
NEWHALL, A. G., 18, 386
Nicandra physaloides, Solanum virus-4 in, 170
Nicotiana, Solanum virus-4 in spp. of, 169–170
tabacum (See Tobacco)
virus-1, combination with Solanum virus-4, 173
Nicotine-oil sprays, compatibility with copper fungicides, 12
NIKITIN, A. A., 19
Nitrogen-deficiency, effect on Phytomonas stewartii, 17
Nitrogen supply, sweet corn, effect on bacterial wilt, 1051
tobacco, effect on mosaic virus concentration, 22
Nodules, legume root-, auxin increase in, 15
Nomenclature, of phytopathogenic bacteria, 128
of viruses, 422, 431
NUGENT, T. J., (5), 19
Nummularia atropunctata, hardwood decay by, 2
Nurseries, black locust, root-knot control by heat, 4
forest, disease developments in, 14
hemlock, rust control, 463
Nutrition, fungus, 592

Oak(s) (See also Quercus)
laurel- (See Quercus laurifolia)
Oats, cold resistance as influenced by crown and stem rusts, 763
crown rust, developmental factors, 659
urediospore viability, 21
variety reactions to, 673
Fusariums in diseases of, 7
loose smut, as influenced by environment, 22

stem nematode disease, incidence and resistant varieties, 827
stem rust, as influenced by barberry eradication, 24
OCFEMIA, GERADO OFFIMARIA, 512
Oidium euonymi-japonica, control by water sprayed under pressure, 289
Oils, animal and vegetable, fungicidal value, 5, 24
Oligostroma acicola, transfer to Scirrhia acicola, 1076
OLIVER, SIDNEY J., (826)
downy mildew, control by spraying vs. eradication of perennial onions, 18
culture trials of fungus, 828
infection in relation to moisture, 933
pink root, host-parasite relations, 629
smut, control by early planting, 6
OORT, A. J. P., 717
Orange, sour-, infectious variegation, 911
Orchards, Armillaria mellea infestation of soils in, carbon bisulphide control, 827
diseases, book review, 915
fundamentals of fruit production, book review, 1002
Ornamentals, diseases and pests, book review, 1002
Orthophosphoric acid, crude, in conifer damping off control, 14
Ortho-toluenesulfonylamide, fungicidal value, 979
ORTON, C. R., 381
OWENS, O. P., (752)

Papaw (See Asimina triloba)
PAPE, HEINRICH, 1002
Para crinkle, potato, on American vs. European varieties, 49
Para rubber tree, Fomes lignosus on, 14
Paradichlorobenzol, for tobacco downy mildew, 216
Para-toluenesulfonylamide, fungicidal value, 979
Parasites, clonal varieties in relation to, 9
PARKER, K. G., (547)
PARKER, R. B., (754)
Paspalum, ergot of spp. (including new hosts), 365
leaf blotch (Ciboria associated) on spp., 749
Pasteurization, of soil, electrical devices for, 18
Pathogens, bacterial phyto-, poured-plate vs. single-cell methods for studying, 852
taxonomy, 128
new etiological terminology, 459
soil, chemical control, 25
Pea, broad-bean mild mosaic infecting, 452
cucumber-type viruses of, 25
damping off, control by cuprous oxides, 306
dieback virus (new), 25
freezing injury to canning-, 188
marsh spot due to manganese deficiency, 3
seedling freezing injury, nature and plant response, 188
Septoria blight, varietal reactions, 25
stem-nematode disease of canning-, incidence and damage, 827
stem-streak virus (new), 25
Peach, Bordeaux as summer fungicide for, 11
brown rot, control, 11
canker, fungi involved, 10
Coryneum blight, control by Bordeaux-petroleum oil emulsion, 828
phony disease, virus, inactivation by heat in nursery trees, 12
localization in woody cylinder, 12
powdery mildew compared to that on rose, 282
scab, control, 11
Sclerotinias of, in Calif., 337
in Wash., 825
spraying by airplane, 823
Peanut, Cercospora leaf spots, fungicidal control, 18, 751
oil, fungicidal value, 5
Pear, fire blight, bacterial survival in dried vs. moist ooze, 142
exudate properties and nature, 142
stony pit of fruits, distribution, symptoms and etiology, 260
transmissibility, 13
Pecan, Phymatotrichum root rot, as influenced by forage and intercrops, 827
pathogenicity and morbid histology, 823
soil treatments for, 827
PECHUMAN, L. L., (547)
PELTIER, GEORGE L., 485
Penicillium, antagonism among spp., 235
corn seedling injury and factors influencing, 231
pines infected by, 1036
oxalicum, substance toxic to corn produced by, 237
Pepper, damping-off, seed and soil treatments for, 750
Verticillium wilt, 359
Peptone broth, growth substances produced by crown gall and related bacteria on, 16
Perennial canker, apple, as influenced by wound dressings, 739
Perkinsiella vastatrix, as vector of Fiji disease of sugar cane, 512
Peronospora destructor, culture trials and factors in, 828
infection by, moisture relations, 933
control by spraying vs. eradication of perennial onions, 18
effusa, infection by, moisture relations, 933
nicotainae, in South America, 291
tabacina, benzol vapor toxicity to, 177
control, by benzol vapors, concentration, 103
by oil-copper combinations, 5
by oil sprays, 24
paradichlorobenzene as eradicant fungicide, 216
endemism, inoculum sources, sporangial dissemination and weather relations, 194

on flue-cured tobacco, epidemic, 79
infection studies, 79
in South America, 291
Persicaria portoricensis, Corticium microsclerotia on, 560
PERSON, L. H., (17) 19, 750
Pestalozzia funerea, pines infected by, 1036
Pests, of ornamentals, book review, 1002
plant, handbook, review, 558
Petroleum oil emulsion, effect on fungicidal value of Bordeaux, 828
Petunia nyctaginiglora, Solanum virus-4 in, 170
Phaseolus, Corticium microsclerotia on spp., 560
Erysiphe polygoni on spp. and vars., 824
acutifolius, powdery mildew resistance, inheritance, 824
coccineus, powdery mildew resistance, inheritance, 824
vulgaris, nodule hyperauxony, 15
Phenothiazine, fungicidal activities of derivatives and, 7, 700, 707
Phenothiazone, fungicidal value, 707
Phoma terrestris, pigment, pH in relation to, 629
isolation and nature, 629
Phomopsis juniperovora, incidence on native red cedars, 991
red cedar nursery stock infection by, source and control, 991
vaccinii, blueberry twig blight due to, 136
Phony disease, peach, virus inactivation by heat in nursery trees, 12
virus localization in woody cylinder, 12
Phycomyces blakesleeana, growth substance relations, 592
Phyllody, in bean varieties, morphology and anatomy, 823
in carrot, 823
in curly-top infected plants, 823
in squash, 823
in tomato varieties, morphology and anatomy, 823
Phymatotrichum omnivorum, control, green manures for, 752
on cotton, as influenced by stem girdling, 753
rainfall relations, 752
distribution and prevalence in Oklahoma, 485
factors influencing spread, 485
as influenced by heavy metals, 2
life history and cytology, unusual features in, 498
on pecan root, as influenced by forage and intercrops, 827
pathogenicity and morbid histology, 823
soil treatment for, 827
Physiologic races, of Actinomyces scabies, 204
determination methods, 609
of Fomes lignosus, 14
of Fusarium vasinfectum, 21
of Puccinia antirrhini, 18
of Puccinia graminis, 22
of Puccinia graminis tritici, 4
of Puccinia rubigo-vera tritici, 4, 962
in Rhizoctonia, 609
of Tilletia levis, 863
of Tilletia tritici, 863
in Uromyces, 9, 820
Phytomonas **aceris**, n. sp., maple leaf spot due to, 968
begoniae, begonia leaf spot due to, 636
flavozonatum, identity with P. begoniae, 636
malvacearum, in irrigated cotton, factors in persistence and distribution, 754
stewartii, as influenced by nitrogen supply in culture, 17
motile bacterium inducing symptoms like those by, 1062
sweet-corn wilt induced by, as influenced by nitrogen supply, 17, 1051
tumefaciens, effect on young apple trees, 21
pathogenicity basis, 16
growth substance production in peptone broth by, 16
PHYTOPATHOLOGY, errata (Vol. 28), 390
Phytopathology, forest-, book review, 915
handbook, review, 558
new etiological terminology, 459
sugar cane diseases, book review, 220
Phytophthora, in infected material, isolation method, 290
cactorum, Lychnis viscaria stem rot due to, 652
strains differing in host pathogenicity, 654
cryptogea, Gloxinia corms, stems and leaves infected by, 17
megasperma, sweetclover root rot due to, breeding and selection for resistance, 909
Pigment, of Phoma terrestris, 629
PINCKARD, J. A., 79, (103), 177, 216, 242, 751
Pine, longleaf (See Pinus palustris)
red (See Pinus resinosa)
western yellow (See Pinus ponderosa)
white (See Pinus strobus)
Pineapple, nematode injury restricted by fungi, 596
yellow-spot disease, distribution and insect vector, 285
Pink root, onion, host-parasite relations, 629
Pinus, bacteria infecting, 1036
distribution of spp. as influenced by damping off vs. soil type, 367
mycorhizas and pseudomycorhizas on spp., effects and factors influencing, 14
wood-staining and -destroying fungi on, relative importance and seasonal prevalence in United States, 1033
palustris, brown-spot needle blight, taxonomy of fungus, 1076
ponderosa, Ceratostomella ips associated with Ips lecontei on, 556
resinosa, mycorhizas and pseudomycorhizas on, environal relations, 24
parasites, environal relations, 24
strobus, Cronartium-induced canker development on young, 1067

nurseries, pH as influenced by alkaline sand and sulphur, 14
PIRONE, P. P., 19, (956)
Pisum sativum (See Pea)
var. arvense, mild mosaic of broad bean infecting, 452
Pityothamnus angustifolius, Corticium microsclerotia on, 561
PLAKIDAS, A. G., 19, 20, 751
Planetree, California (See Platanus racemosa)
Plants, diseases, pests and protection, handbook, review, 558
insect populations of individual, determination method, 20
Plasmodiophora brassicae, crucifer infection as influenced by soil amendments, 9
Plasmopara viticola, mycelium overwintering in wild grape, 822
Platanus racemosa, leaf spot fungi of, sporulation induced by host-juice media, 821
Pleospora lycopersici, inoculations on tomato and legumes, 541
relationship to Macrosporium sarcinaeforme, 541
Poison-hemlock ringspot, aphid transmission, 824
POLHAMUS, LOREN G., (970)
Polyporus glomeratus, cankers on maple and beech due to, 3
sterile conks and associated cankers on beech, 3
schweinitzii, on red pine, 24
POOLE, R. F., 750
POPHAM, W. L., (22)
Populations, insect, determination method for individual plants, 20
Populus, Nectria cankers, avenues of entry, 353
PORTE, W. S., 757
Portraits, Meier, Fred Campbell, 302 and facing p. 293
Potato, blackleg, bacteria associated, 831
Bliss Triumph var., differential host of tomato tip blight, 162
blue-stem, suberin and suberized deposits in, 278
Canada streak, distribution, 932
Lygus-induced wilt, non-identity with purple-top, 14
nematode infestation of roots, 287
net necrosis, suberin and suberized deposits in, 280
purple-top disease, etiology and insect transmission, 14
Rhizoctonia solani penetration of tubers, seed treatment relations, 759
scab, control by calomel, 13
as influenced by soil pH, 750
isolation method for pathogen, 1000
resistance in relation to pathogenic races of Actinomyces, 204
scurf, control by calomel, 13
seed-piece decay, bacteria associated, 831
spindling sprout, non-transmissible form, 21
suberin and suberized deposits in normal vs. diseased, properties, 274
tuber blotch, distribution, 932
virus-B (See Solanum virus-4)
viruses affecting European vs. American, 40
X-virus, hosts, 57
immunity inheritance, 362
inhibition by insect juices, 323
yellow-dwarf virus, as influenced by insect juices, 2, 321, 322, 323
yellow mosaics, 6
of European and American varieties, comparisons and classification of viruses, 917
Powdery mildew (See under Mildew)
Pox, sweet-potato, Actinomycete strains in relation to, 17
Preservatives, wood-, acceleration of toximetric tests by soil, 901
PRESLEY, JOHN T., (752), 498
PRICE, W. C., 20, 83, 903
PROEBSTING, E. L., (10)
Propagation, by clonal varieties, diseases in relation to, 9
Protein, tobacco-mosaic virus-, molecular weight, 17, 495
purification and properties, 250
X-ray diffraction, 17
Prune, drought spot, as boron deficiency, 650
Sclerotina spp. on, in Wash., 825
Taeniothrips inconsequens population in orchards, as influenced by calcium cyanamid, 825
Prunus, Sclerotinias on spp., apothecia destroyed by calcium cyanamid, 436
in Calif., 337
control, 11
Tranzschelia pruni-spinosae on native Calif. spp. 645
pennsylvanica, Nectria cankers, avenues of entry, 353
serotina, Tranzschelia pruni-spinosae on, germination of overwintered teliospores on fallen leaves, 74
PRYOR, DEAN E., 641
Pseudo-calico, as celery virosis (new), and host range, 824
Pseudomycorhizas, on pines, 14
on red pine, environal relations, 24
Pseudo-net-necrosis, potato, 6
differentiation and classification, 925
geographical distribution, 932
Pseudoperonospora cubensis, control by copper fungicides, 751
infection in relation to moisture, 933
humuli, infection in relation to moisture, 933
Psorosis, citrus, infectious variegation in relation to, 911
registration of trees inspected for, 825
relation to other virus-like effects on, 6
Puccinia, urediospores, light effects on germ tubes of spp., 667
antirrhini, infection in relation to moisture, 933
snapdragons resistant to two races of, 18

coronata, developmental factors, 659
effect on cold resistance of oat vars., 763
oat variety reactions to, 673
avenae, urediospore viability, factors influencing, 21
graminis, barley variety resistant to, 209
epidemiology in Mexico, 22
incidence as influenced by barberry eradication, 24
physiologic races in Mexico, 22
avenae, effect on cold resistance of oat varieties, 763
as influenced by environment, 22
tritici, control by toluene compounds, 978
Early Blackhull wheat resistant to, 644
physiologic races, temperature effects on infection and development, 4
temperature effects on urediospore germination and germ tube development, 4
helianthi, infection in relation to moisture, 933
rubigo-vera tritici, Agropyron trichophorum infected by, 965
physiologic races in 1938 epidemic, 962
wheat hybrids resistant to, 965
triticina, inoculum source for fall infections, 4
urediospores air-borne, 4
Pullularia pullulans, pines infected by spp., 1036
Purple-top disease, potato, etiology and insect transmission, 14
Pythium, damping-off in eggplants, peppers and tomatoes due to, 750
incidence of spp. on cotton, 754
Persian buttercup root rot due to spp., 828
watermelon blossom-end rot due to spp., 391
acanthicum, parasitism and general study, 395
anadrum, parasitism and general study, 415
aphanidermatum, cucurbit root and fruit rot due to, 624
growth, reproduction and hosts, 624
helicoides, parasitism and general study, 408
mastophorum, morphology, development and relationships, 1005
megalacanthum, morphology, development and relationships, 1005
periplocum, parasitism and general study, 402
polymastum, morphology, development and relationships, 1005
ultimum, control by cuprous oxides, 306

Quasi-factorial design of experiments, randomized-block design compared, 637
Quercus, Cronartium spp. on Castanea, Castanopsis, Lithocarpus and, 998
decay due to Ustulina in living, 2
heart rot due to Stereum gausapatum, 504
Nectaria cankers, avenues of entry, 353
virus-like symptoms on, 8
laurifolia, citrus-leprosis-like lesions on, 907
Quince, rust, comparison with other rusts affecting apple, 801
QUIRK, AGNES J., 20

RAEDER, J. M., 1001
Rainfall, effect on cotton root rot, 752
Randomized-block design of experiments, quasi-factorial design compared, 637
Ranunculus asiaticus, root rot due to Pythium spp. in, 828
Rapeseed oil, fungicidal value, 5
Raspberries, black, mosaic, transmission by Amphorophora, 825
mosaic-resistant, breeding and factors in, 647
mosaic-vector resistant, breeding and factors in, 647
RAWLINS, T. E., 20
Red cedar (See Juniperus virginiana)
Red clover (See Trifolium pratense)
Red leaf disease, of grapes, control by mite control, 10
REDDICK, DONALD, 384, (385)
REDDY, C. S., 382
REEVES, E. L., 739
REINKING, OTTO A., 558
Report, American Phytopathological Society, annual meeting (1938), 375
summer meeting, 1003
Resistance (See also Susceptibility)
barley, to various diseases, 209
beans, to anthracnose, inheritance, 1
to powdery mildew, 824
blueberry, to powdery mildew, 545
cereals, to rusts, 672
cherry, to buckskin disease, 20
cotton, to wilt, 753
to wilt/nematode complex, 755, 974
cowpea, to diseases, breeding for, 826
to powdery mildew, 824
cucumber, to bacterial wilt, 996
currant, to blister rust, 26, 981
mechanism, 38
disease-, clonal varieties in relation to, 9
time-weighting method of comparing, 873
oats, to cold, as influenced by rusts, 763
to stem nematode, 827
pea, to Septoria blight, 25
potato, to scab, 204
to X-virus, inheritance, 362
raspberry, to mosaic, 647
snapdragon, to rust, 18
sugar beet, to curly top, virus concentration in relation to, 825
sweetclover, to root rot, 909
time-weighting method of comparing, 873
tomato, to diseases, Lycopersicum chilense as source, 215
to Fusarium wilt, 24, 25, 757, 871
study technique, 945

to leaf mold, 9
to leaf spots, 750
watermelon, to wilt, 5
wheat, to bunt, 863
to leaf rust, 965
wheat variety, to stem rust, 644
Reviews, biometry methods, 657
book, bacteriology manual, 128
diseases of fruits and hops, 915
diseases of ornamentals, 1002
fruit production fundamentals, 1002
fungi of India, 374
phytopathology handbook, 558
sugar-cane diseases, 220
statistical methods, 657
Rhizobium phaseoli, auxin increase in nodules induced by, 15
Rhizoctonia, damping off in eggplants, peppers and tomatoes due to, 750
potato scurf due to, control by calomel, 13
bataticola, cowpea breeding for resistance, 826
incidence on cotton, 754
microsclerotia, bean web blight due to, 559
solani, bean stem rot due to, host-parasite relations as influenced by nutrients, 1
cultural characters and pathogenicity, 793
culture variant in physiology and pathogenicity, 267
incidence on cotton, 754
perfect stage from cotton seedlings, 373
physiologic races, determination methods, 609
potato-tuber penetration, seed-treatment relations, 759
strains causing crown rot vs. dry-rot canker of sugar beets, 796
sugar beet dry-rot canker due to, 793
RHOADS, A. S., (907)
Rhododendron, bud and twig blight (Sporocybe azaleae), 517
Rhopalosiphum melliferum, as vector of celery yellow-spot virus, 824
Rhus, Corticium galactinum infecting, 5
Ribes (See also Currant)
alpinum, Cronartium ribicola immunity of a staminate clone, 981
punctatum, Cronartium ribicola susceptibility, 643
Rice, leaf blotch (new), Ciboria associated, 749
RIKER, A. J., (16), 21, 382, 852
Ringspot, poison-hemlock, aphid transmission, 824
tobacco No. 1, virus inhibition by insect juices, 323
virus induced, in China, 461
ROBERT, ALICE L., (6), (284)
ROBERTS, JOHN W., 915
Robinia pseudoacacia, emergence, as influenced by formaldehyde, 14
leaf spots, 14
root-knot control by heat in nursery, 4
systemic brooming virus-induced, 8
winter storage losses, 14
RODENHISER, H. A., (378)
Root-knot, in black locust nurseries, control by heat, 4
chemical soil treatment for, 25
on cotton, in Tenn., 751
varietal susceptibility to Fusarium wilt plus, 974
on cowpea, breeding for resistance to, 826
hosts, incidence and control in California, 752
on tobacco, as influenced by transplanting dates, 21
on tomato, greenhouse control, 11
in Tenn., 751
Root nodules, legume-, auxin increase in, 15
Root rot(s) (See also Rots)
apple white-, Corticium causing, 5
biological control, 750
black, of flue-cured tobacco, factors influencing, 751
of Persian buttercup, Pythium spp. causing, 828
Phymatotrichum, control, 752
as influenced by heavy metals, 2
life history and cytology, 498
survey and factors influencing, Oklahoma, 485
on cotton, as influenced by stem girdling, 753
rainfall relations, 752
on pecan, 823
as influenced by forage and intercrops, 827
soil treatments for, 827
Pythium aphanidermatum causing, 624
soil treatment for, 750
sweetclover Phytophthora-, breeding and selection for resistance, 909
wheat, seed treatments for, 16
Rose, black spot, control by cuprous oxides, 315
powdery mildew, compared to that on peach, 282
control by water sprayed under pressure, 289
ROSEN, H. R., 21
Rosette, virus-induced, in China, 460
Rotations, crop, in nematode control, 752
sweet-potato diseases in relation to, 750
Rot(s) (See also Root rots)
bean stem-, Rhizoctonia-host relations as influenced by nutrients, 1
butt- and trunk-, Daedalia causing in hardwoods, 3
cotton boll-, fungi associated, 753
cotton internal collar-, Thielaviopsis causing, 751
cowpea charcoal-, breeding for resistance, 826
cucurbit Pythium root- and fruit-, 624
of forest trees, book review, 915
fruit- and root-, Pythium aphanidermatum causing, 624
of hardwoods, by Ustulina and other Ascomycetes, 2

Kale black-, seed treatment by chloropierin, 19
Lychnis stem-, Phytophthora cactorum causing, 652
narcissus bulb-, mercuric oxide soil treatment for, 23
oak heart-, Stereum gausapatum causing, 504
potato seed-piece, bacteria associated, 831
sugar beet crown-, 796
sugar beet dry-rot canker, 793
sweet potato soil-, control, 750
watermelon blossom-end, Pythium spp. causing, 391
tomato blossom-end, environal/metabolic factors, 7
wood, testing of preservatives against, 901
Rubber tree, Para, Fomes lignosus on, 14
Rubus, Corticium galactinum infecting spp., 5
stamen blight of spp. due to Hapalosphaeria deformans, 829
cuneifolius, Corticium microsclerotia on, 561
leucodermis, mosaic, aphid transmission, 825
RUDOLPH, B. A., (359), 753
RUE, JOHN L., (12)
Rugose mosaic, virus-induced, in China, 461
Rust(s), apple, pathogenicity, symptoms and 3 fungi compared, 801
varietal susceptibility in relation to spp., 18
barley stem-, variety resistant to, 209
bean, infection in relation to moisture, 933
physiologic races, 9, 820
cereal, incidence of grasses in relation to, 24
nature of resistance to, 672
cereal stem-, as influenced by barberry eradication, 24
infection in relation to moisture, 933
on Jasminum spp., 783
oats crown-, developmental factors, 659
effect on cold resistance, 763
urediospore viability, 21
oat stem-, effect on cold resistance, 763
pine-oak, occurrence on Castanea, Castanopsis and Lithocarpus, 998
on Prunus spp. native to California, 645
quince, comparison with other rusts affecting apple, 801
red cedar, infection time and control as influenced by aeciospore germinability and dissemination, 812
snapdragon, resistant lines, 18
spinach white-, 616
epidemic, 12
wheat leaf-, inoculum source for fall infections, 4
physiologic races in 1938 epidemic, 962
resistant hybrids, 965
wheat stem-, control by toluene compounds, 978
Early Blackhull wheat resistant to, 644
epidemiology in Mexico, 22
as influenced by barberry eradication, 24
physiologic races in Mexico, 22
temperature effects on infection and development of physiologic races, 4
white, of spinach, 12
white pine blister- (See Cronartium ribicola)
Rye, Fusarium spp. in disease of, 7
smut spp. infecting, 576
stem rust, as influenced by barberry eradication, 24
RYKER, T. C., 749

SALMON, S. C., 657
Salpiglossis variabilis, Solanum virus-4 in, 170
Sambucus sampsonii, Corticium microsclerotia on, 561
Saperda tridentata, as vector of Dutch elm disease, 547
Saprogen, new term, 459
Saprogenic, new term, 459
Saprogenicity, new term, 459
Saururus cernuus, Corticium microsclerotia on, 561
Scab, apple, control by phenothiazine and derivatives, 7, 707
barley, variety resistant to, 210
peach, control, 11
potato, control by calomel, 13
as influenced by soil pH, 750
isolation method for pathogen, 1000
resistance in relation to pathogenic races of Actinomyces, 204
SCHAAL, L. A., 759
SCHROEDER, F. R., (485)
SCHULTZ, E. S., 21, (362)
SCHWARTZE, C. D., 647
Scirrhia **acicola**, n. comb., perfect stage of pine brown-spot needle-blight fungus, 1076
Sclerotinia, ascospore discharge, method for, 373
camellia blossom blight due to sp., 824
fructicola, apothecia, calcium cyanamid destroying, 436, 825
in peach canker complex, 10
spores, toxicity of Bordeaux vs. oil-Bordeaux to, 828
on stone fruits, in Calif., 337
in Wash., 825
thrips transmitting, 825
fructigena, fungicidal action of phenothiazine and derivatives on, 701
laxa, on stone fruits, in Calif., 337
in Wash., 825
Sclerotium bataticola, conifer seedling infection, 14
rolfsii, incidence on cotton, 754
indexing for infection on sugar beets, and factors influencing, 15
Scolytus, Ceratostomella ulmi fructifications in galleries of spp., 551
Scorch, strawberry, new Mycosphaerella leaf spot resembling, 20
SCOTT, C. E., 752

SCOTT, G. W., (820)
Scurf, potato, control by calomel, 13
Sea spray, effect on cypress canker fungus, 823
Seed treatment(s), for bean bacterial blight, 19
cereals, disinfectants compared, 16
chlorine gas as disinfectant, 913
cotton, 754
effect on: permeability of testa, 753
seedling survival, 755
as influenced by delinting, 754, 755
by organic mercury dust, and H_2SO_4, 755
as influenced by type and period of storage, 754, 986
by sulphuric acid, 879
for damping off, 750
kale, for black rot, 19
measuring interfacial friction of dusted seeds, 200
potato, fungus-tuber penetration in relation to, 759
sugar beet, factors influencing, 15
wheat seedling blight, 16
Seed-corn maggot, as vector of Erwinia carotovora, 849
Seeds, diseases borne by, in cereals, 8
chlorine gas disinfection for, 913
sweetclover fungi, 912
broad-bean mild mosaic, 454
dusted, measuring interfacial friction, 200
testa, permeability in cotton as influenced by seed treatment, 753
weed, electrical pasteurization of soil to kill, 18
Selocide, mite and red-leaf control on grapes by, 10
Septoria, tomato leaf spot due to, varietal reactions, 750
apii, dissemination on clothing, 553
pycnidial numbers per leaf spot, 646
spore numbers per pycnidium, 646
bromigena, Bromus inermis leaf spot due to, 1
susceptibility of inbred lines, 554
sector variants, 554
variability vs. stability of monosporous strains, 554
pisi, overwintering, 25
varietal reactions to, 25
Setaria viridis, corn crazy-top-like symptoms in, 820
Sesame oil, fungicidal value, 5
SEVERIN, HENRY H. P., (824), 826
SHANDS, R. G., 209
SHAW, K. J., 21
SHAW, LUTHER, (79), 382, 751
Shelterbelt plantings, Phymatotrichum root rot in relation to, 485
SHERBAKOFF, C. D., 21, 751
Shrubs, ornamental, diseases and pests, book review, 1002
SIGGERS, PAUL V., 1076
Single-cell culture technique, poured-plate method compared, 852
Sinningia speciosa, Phytophthora cryptogea infecting corms, stems, and leaves, 17
Sitanion jubatum, Tilletia spp. on, 577
Ustilago spp. on, 490
SLOWATA, STANLEY S., (529)
Small fruits, diseases, book review, 915
SMITH, A. L., 756
SMITH, CLAYTON O., 645, (821)
SMITH, DONALD J., 821
SMITH, T. E., 22
Smut(s) (See also Bunt)
barley covered-, as influenced by environment, 22
barley loose-, infection factors, 717
cereal-, chlorine gas seed disinfection for, 913
corn, growth types and factors influencing, 473
forage-grass susceptibility to spp., 576, 490
oats loose-, as influenced by environment, 22
onion, control by early planting, 6
wheat loose-, infection factors, 717
Snapdragon, Botrytis blight, trichomes as infection courts, 651
rust, infection in relation to moisture, 933
resistant lines, 18
SNYDER, W. C., 359, 827
Sodium carbonate, spray residue removal by, 12
Sodium chloride, effect on cypress canker fungus, 823
Sodium hydroxide, in spray residue removal, 12
Soft rots, of potato tubers, bacteria associated, 831
Soil(s), carbon dioxide evolution from, relation to black root rot of flue-cured tobacco, 751
disinfection, by calcium cyanamid, 436
fungus contamination technique, 755
hydrogen-ion concentration, effect on plant pathogens, 9, 750, 756
nematode populations, 601
orchard, Armillaria mellea control by carbon bisulphide in, 827
pasteurization, electrical devices for, 18
pathogens in, chemical control, 25
pine distribution in relation to, 367
-rot, of sweet potato, control, 750
sweet potato diseases persisting in, 750
treatments, for damping off, 750
for root rot fungi, 750
weed seeds in, chemical control, 25
wood preservatives tested in, 901
Solanum, Solanum virus-4 in spp., 59, 170, 171
capsicastrum, differential host of tomato tip blight, 162
Solanum virus-2, combination with virus-4, 173
Solanum virus-4, on American vs. European potato varieties, 59
combination with other viruses, 173
host range and differentiation, 169
separation from virus-1, 168
Solidago, Elsinoë solidaginis on new host spp. of, 970
Sophora, virus-like symptoms on, 8

Sorghum, covered smut, chlorine gas seed disinfection for, 913
SPAULDING, PERLEY, (351)
SPENCER, ERNEST L., (17), 22, 502, (1051)
Sperm oil, fungicidal value, 5
Spaceloma **lippiae**, n. sp., Lippia lanceolata anthracnose due to, 655
menthae, comparison to S. lanceolata, 654
Sphaelotheca sorghi, lysis in crosses of, 14
pannosa, control by water sprayed under pressure, 289
forms on peach vs. rose compared, 282
Spinach, downy mildew, infection in relation to moisture, 933
Spinach, white rust (Albugo occidentalis), 616
epidemic, 12
Spindle tuber, virus induced, in China, 460
Spindling sprout, potato, non-transmissible form, 21
Spores, fungus, germination trials, and factors influencing, 828
with fungicides, 16
isolator for, 212
Sporocybe azaleae, azalea and rhododendron bud and twig blight due to, 517
Sporulation, of fungi, culture media for, 821
Spot necrosis, virus-induced, in China, 461
Spray injury, by copper fungicides as influenced by oil supplements, 24
Spray residue, removal methods, 12
Spraying, by airplane, in peach orchards, 823
Sprays, bentonite sulphur, colloidality, 8
liquid lime-sulphur, standardized procedures important in diluting, 993
Squash(es), Choanephora cucurbitarum attacking, 898
crookneck, Pythium root and fruit rot of, 624
phyllody in, 823
STAKMAN, E. C., 22
Stamens, blight of blackberry-, Hapalosphaeria deformans causing, 829
STANDEN, J. H., 656
STARK, F. L., (11)
Statistical analysis, of replicated experiments, review, 657
Stem-end browning, potato, suberin and suberized deposits in, 278
Stem rot, bean Rhizoctonia-, host-parasite relations as influenced by nutrients, 1
of Lychnis viscaria, Phytophthora cactorum causing, 652
Stem rust, wheat, control by toluene compounds, 978
Stem streak virus of pea, 25
Stereum gausapatum, oak heart rot due to, 504
growth and variability in culture, 504
STEVENS, H. E., 537
STEVENS, NEIL, 385
STEVENSON, F. J., 362
STEVENSON, JOHN A., 374, 915, (616)
Stigmella platani-racemosae, sporulation induced by host-juice media, 821
Stigmina platani, sporulation induced by host-juice media, 821
Stipple streak, potato, on American vs. European varieties, 51
Stone fruits, brown rot, apothecia destroyed by calcium cyanamid, 436
control, 11
distribution in Calif., 346
Sclerotinias of, in Calif., 337
in Wash., 825
Nectria cankers, avenues of entry, 353
thrips population in orchards as influenced by calcium cyanamid, 825
Tranzschelia pruni-spinosae infection, germination tests, 72
on native Calif., spp., 645
Stony pit, pear, distribution, symptoms and virus etiology, 260
transmissibility, 13
Storage, effect on mercury-dust-treated cotton seed, 986
Strawberry, Cercospora leaf spot, (apparently new), 19
Mycosphaerella leaf spot (new), 20
scorch, new Mycosphaerella leaf spot resembling, 20
Streak virus, in China, 460, 461
"up-to-date," (See Solanum virus-4) 168
STREETS, R. B., 827
Stripe disease, barley variety resistant to, 211
Strumella coryneoidea, hardwood decay by, 2
Suberin, microchemical and optical properties in normal vs. diseased potato tubers, 274
Suberized deposits, microchemical and optical properties in normal vs. diseased potato tubers, 274
Sugar beet, crown rot, comparison to dry-rot canker, 796
curly top virus-1, concentration in root tips of resistant vs. susceptible vars., 825
damping-off, quasi-factorial vs. randomized-block designs of experiments on, 637
seed treatments and factors influencing, 15
dry-rot canker, 793
Rhizoctonia solani culture variant from, 267
Sclerotium rolfsii infection, indexing for and factors influencing, 15
seed treatment, factors influencing, 15
Sugar cane, anatomy, in book review, 220
diseases, book review, 220
Fiji disease, behavior of POJ 2878 to, 512
immunity reactions in testing virus classifications, 7
SUIT, F. R., 200
Sulphur, effect on pH in white-pine nurseries, 14
elementary fungicidal, particle size, 8
fused bentonite-, colloidality, 8
in sweet potato soil rot control, 750
Sulphuric acid, cotton-seed treatment by, effect on fungi and bacteria, 879
Sumac, Corticium galactinum infecting, 5
Sun-blotch, avocado, in Florida, 537

Sunflower, rust, infection in relation to moisture, 933
Susceptibility (See also Resistance)
apple, to rusts, 18
Bromus inermis, to Septoria leaf spot, 1, 554
currant, to blister rust, 643
grasses, to cereal smuts, 490
Sweetclover, white (See Melilotus alba)
Sweetclovers, fungus parasites infecting seed of, 912
Sweet potato, Actinomycete strains compared for pathogenicity on, 17
diseases, crop rotation in relation to, 750
pox, etiology, 17
soil rot, control, 750
Syntherisma villosum, Corticium microsclerotia on, 561

Taeniothrips inconsequens, population in prune orchards as influenced by calcium cyanamid, 825
TAPKE, V. F., 22
Taxonomy, of Fusarium group, factors influencing identification, 827
of phytopathogenic bacteria, 128
of viruses, 422, 431
TAYLOR, A. L., 752
Technique (See also Culture media)
analysis of variance, review, 657
for ascospore discharge in Sclerotinia, 373
benzol vapor concentration determination, 179
chlorine gas disinfection of seed, apparatus for, 914
culture, Fusarium identification and classification in relation to, 827
poured-plate efficacy for bacterial plant pathogens, 852
renewed liquid, for fungi, 895
transfer hood, 908
culture-tube label, 761
experimental, quasi-factorial vs. randomized-block designs, 637
fungicides, adherence properties, determination by cataphoresis, 19
toxicity test method for volatile, 641
for fungus soil contamination, 755
fungus spore germination trials and factors in, 828
generalized standard errors in wheat bunt tests, review, 657
for host resistance and pathogenicity studies, 945
inoculation, for fungi, water-culture method, 23
for Fusarium vasinfectum, 755
insect populations of individual plants, determination, 20
insect vector inoculation chamber, 649
isolation, for Actinomycetes, 1000
for Phytophthora spp., 290
single-cell, efficacy for bacteria, 852
measuring interfacial friction of dusted seeds, 200
for nematode infestation estimation, 287
physiological races, determination, 609
quasi-factorial vs. randomized-block designs of experiments on damping off, 637
resistance to disease, time-weighting method of comparing, 873
for soil contamination with Fusarium vasinfectum, 756
spore isolator, 212
statistical analysis of replicated experiments, review, 657
for suberin vs. suberized deposit determinations, 275
toxicity testing of volatile compounds, 641
toximetric tests of wood preservatives, acceleration by soil, 901
Temperature, effect on viruses, 20, 245, 441
Terminology, etiological, new, 459
Test tubes, label for, 761
Tetranychus pacificus, on grapes, control by Selocide, 10
rôle in red-leaf disease, 10
Thallium injury, to tobacco, frenching in relation to, 502
Thamnotettix geminatus, as vector of aster yellows in delphiniums, 826
montanus, as vector of aster yellows in delphiniums, 826
THARP, W. H., 756
Thelephora galactina (See Corticium galactinum)
Thermal "death" point, of tobacco viruses, 20, 245
Thiamin, Phycomyces growth in relation to, 592
Thielaviopsis basicola, cotton internal collar rot apparently due to, 751
in flue-cured tobacco, factors influencing, 751
as influenced by soil pH, 750
Thionol, fungicidal value, 707
THIRUMALACHAR, M. J., 783
THOMAS, HAROLD E., 827
THOMAS, H. EARL, (824)
THORNBERRY, H. H., 250
Thrips, population in prune orchards as influenced by calcium cyanamid, 825
as vectors of Sclerotinia fructicola, 825
tabaci, as vector of tomato tip-blight virus, 159
THURSTON, H. W., Jr., 993
Tilia americana, Nectria canker, growth and mortality relations, 23
Tilletia levis, forage grass susceptibility to, 575
perennial mycelium of, 585
purified physiologic races, reinoculation of resistant wheat varieties with, 863
wheat chlorotic mottling due to, 456
tritici, forage grass susceptibility to, 575
purified physiologic races, reinoculation of resistant wheat varieties with, 863
Tip blight, tomato, virus (new) causing, 156
TISDALE, W. B., (382)
Tobacco, Bacterium angulatum infection as influenced by water-soaking, 2, 889
Bacterium tabacum infection as influenced by water-soaking, 889

benzol vapor toxicity to, 186
black root rot of flue-cured, CO_2 evolution from soils in relation to, 751
curly top infection, recovery and acquired tolerance, 743
of Turkish, antibody-like substance formed, 828
differential host of tomato tip blight, 164
downy mildew, control, 198
by benzol vapors, concentration, 103, 177
by oil-copper combinations, 5
by paradichlorobenzine as eradicant, 216
endemism, inoculum sources, dissemination and weather relations, 194
of flue-cured, infection studies, 79
species (2) reported from South America, 291
flue-curing, temperatures, 246
effect on infectivity of mosaic virus-1, 242
frenching, thallium injury in relation to, 502
genotypes, reactions to cucumber and potato viruses, 16
Granville wilt, control by rotation, 22
weeds as reservoirs of, 22
leaves, injury by water-soaking, 884
mosaic, temperature relations of virus and host, 441
virus, concentration as influenced by nitrogen nutrition, 22
detection in plant by rapid reagent-indicator method, 20
inhibition by insect juices, 2, 321
molecular weight of protein of, 17, 495
movement, in tissues and factors influencing, 1
in tomato plants, 684
physico-chemical study of protein of, 7
thermal inactivation, 20, 245
virus-1, infectivity as influenced by flue-curing, 242
resistant genotypes compared, 16
virus-6, purification and properties of, 250
X-ray diffraction and molecular weight of virus protein prepared by Na_2SO_4 method, 17
necrosis, characteristic macromolecular substances in, 86
ultrafiltration of juices of plants with, 83
virus, inhibition by insect juices, 323
properties, 84
nematodes on, as influenced by transplanting dates, 21
pea viruses infecting, 25
ring-spot virus-1 inhibited by insect juices, 323
seedlings, benzol vapor toxicity to, 177
Solanum virus-4 in, 169
viruses, in China, 461
thermal inactivation of necrosis vs. mosaic, 20
yellow ring spot, symptoms, 549
virus "longevity" in seed, 549
Toluene compounds, plant disease control by, 978
Tomato, blossom-end rot, environal/metabolic factors in development, 7
Chilean, as disease resistance source, 215
Cladosporium leaf mold, resistant red variety, 9
damping off, seed and soil treatments for, 750
disease resistance, Lycopersicum chilense as source of, 215
early blight, control by cuprous oxides, 315
delayed spraying for, 11
Fusarium wilt, chemical soil treatment for, 25
cultural characters and pathogenicity of strains, 24
host resistance and pathogenicity, study technique, 945
resistance to, as influenced by nematodes, 871
time-weighting method of comparing, 873
varietal resistance, 24, 25
tests and factors influencing, 871
hybrid with Lycopersicon hirsutum mosaic-tolerant and wilt-resistant, 757
Macrosporium leaf spot, varietal reactions, 750
phyllody in varieties, morphology and anatomy, 823
root-knot nematodes on, control in greenhouse, 11
in Tenn., 751
Septoria leaf spot, varietal reactions, 750
Solanum virus-4 infecting, 170
susceptibility to Pleospora lycopersici vs. Macrosporium sarcinaeforme, 541
tip blight, differential hosts, 161
insect transmission, 159
virus (new) causing, 156
viruses, in China, 461
TOMPKINS, C. M., (633), 828
TOOLE, E. RICHARD, (463)
Top necrosis, potato, on American vs. European varieties, 61
Torula, hardwoods stained by spp. of, 1037
Toxicity, of thallium, to tobacco, 502
of volatile compounds, testing method, 641
of wood preservatives, testing, 901
Transfer hood, for cultures, 908
Tranzschelia pruni-spinosae, morphology of promycelia, basidia and basidiospores, 74
on Prunus spp. native to Calif., 645
teliospores, germination of overwintered, 72
Trees, clonal varieties for planting, parasites in relation to, 9
diseases, book review, 915
forest, disease developments in nurseries, 14
hardwood, butt and trunk rot due to Daedalia, 3

decay by Ustulina, etc., 2
Nectria cankers on, avenues of entry, 351
insect populations of, determination method, 20
nursery, root-knot on black locust controlled by heat, 4
shade, diseases and pests, book review, 1002
virus and virus-like diseases, 8
wood-destroying and -staining fungi on, 1031
Trichoderma, pines infected by, 1036
Trichomes, as infection courts in Botrytis blight of snapdragon, 651
Trifolium, mild mosaic of broad bean infecting spp., 452
susceptibility of spp. to Pleospora lycopersici vs. Macrosporium sarcinaeforme, 541
pratense, rust, infection in relation to moisture, 933
Tripsacum dactyloides, Aplanobacter stewarti host, 6, 284
TRUE, R. P., 529
Trunk rot, of hardwoods, Daedalia causing, 3
Tsuga canadensis, rust due to Melampsora farlowii and control, 463
caroliniana, rust due to Melampsora farlowii and control, 463
Tuber blotch, potato, 6
differentiation and classification, 925
geographical distribution, 932
Tubes, culture-, label for, 761
TUCKER, C. M., (17)
Tulip, "breaking," Easter lily mosaic virus component causing, 3
first record (?), 1001
mosaic, Easter lily mosaic virus component causing, 3
Tumors, plant (See also Crown gall)
as influenced by colchicine, 221
Tung oil, fungicidal value, 5
Turnip, mosaic, virus inhibition by insect juices, 323
Twig blight, azalea and rhododendron, Sporocybe azaleae causing, 517
blueberry Phomopsis-, 136
hemlock, Melampsora farlowii causing, 463
Tylenchus dipsaci, oats and peas infested by, 827
TYLER, LEON J., 547
Typhlocyba pomaria, Entomophthora sp. from, 9

ULLSTRUP, ARNOLD J., 373, 905
Ulmus, butt and trunk rot due to Daedalia, 3
Cephalosporium die-back, 371
Ceratostomella ulmi, effects on spp., 23
fructifications in England, 551
Dutch elm disease, scouting and sampling as aid to eradication, 529
americana, Dutch elm disease in relation to common elm borer, 547
Ultrafiltration, of tobacco-necrosis plant juices, 83
Urocystis cepulae, onion smut control by early planting, 6
Uromyces fallens, infection in relation to moisture, 933
hobsoni, Jasminum spp. infected by, 783
life history and morphology, 783
phaseoli, infection in relation to moisture, 933
typica, physiologic races, 9, 820
Ustilago, inoculation technique, 717
hordei, chlorine-gas seed disinfection for, 913
forage-grass susceptibility to, 490
as influenced by environment, 22
nigra, forage-grass susceptibility to, 490
nuda, infection factors, 717
tritici, infection factors, 717
zeae, growth types, genetic and environal factors, 473
Ustulina vulgaris, hardwoods rotted by, 2

Vaccinium, blueberry spp., powdery mildew and varietal reactions to, 545
corymbosum, Phomopsis twig blight, 136
VALLEAU, W. D., 549, 884
Valsa cincta, in peach canker complex, 10
leucostoma, in peach canker complex, 10
Variance, analysis of, review, 657
Variegation, infectious, in citrus, 911
Varieties, clonal, diseases in relation to, 9
Vectors (See Insects as vectors)
Vegetables, Pythium infections, 624
Veinbanding, potato, on American vs. European varieties, 51
Venturia inaequalis, control by phenothiazine and derivatives, 7, 707
Vernonia gigantea, Corticium microsclerotia on, 561
VERRALL, A. F., 1031
Verticillium albo-atrum, on cotton, breeding for resistance, 753
control by soil fungicides, 753
pepper wilt due to, 361
Vicia, mild mosaic of broad bean infecting spp., 452
faba, mild-mosaic virus (new) of, 448
host range, transmission, etc., 448
Vigna, Erysiphe polygoni on spp. and vars., 824
sesquipedalis, powdery mildew resistance inheritance, 824
sinensis, Choanephora cucurbitarum on, 898
Corticium microsclerotia on, 561
powdery mildew resistance inheritance, 824
resistance to diseases, breeding for, 826
VINSON, CARL G., (17)
Virulence, bacterial, basis in Phytomonas tumefaciens, 16
as influenced by nitrogen supply, 1051
fungus, as influenced by culture media, 757
Virus disease(s), annotated list (China), 459
of plants, first record (?), 1001
of trees, 8

Virus(es), antibody-like substance to, 828
aster-yellows, 14
in delphinium, 826
avocado, sun-blotch (?), 529
bean, pea viruses in, 25
black locust, systemic brooming due to, 8
black raspberry, mosaic, 825
broad bean, mosaic (new), 448
celery, 824
cherry, buckskin disease, 20
new bud-transmissible (?), 821
citrus, infectious variegation, 911
leprosis (?), 907
psorosis, 6, 825
classification, 422, 431
sugar-cane immunity reactions in testing, 7
cucumber, 16
mosaics, 3, 903
cucumber-type, on pea, 25
currant (new), 369
detection in plant by rapid reagent-/ indicator method, 20
Easter lily, mosaic, 3
grape, new transmissible disease (?), 10
infectivity as influenced by insect juices, 2
inhibition of activity by insect juices, 321
insects as vectors, 2, 14, 159, 286, 321, 450, 512, 647, 824, 825, 826
movements within plant, 684
narcissus, white streak (new), 826, 890
nomenclature, 422, 431
oak, leprosis (?), 907
virus-like symptoms, 8
pea, dieback, 25
stem-streak, 25
peach, phony disease, 12
pear, stony pit (new), 13, 260
pineapple, 285
plant, annotated list (China), 459
immunization to, 828
nomenclature, 422, 431
poison hemlock, ringspot, 824
potato, 16, 40
purple-top, 14
X-virus, 57, 323, 362
yellow dwarf, 2
yellow mosaics, 6, 917
raspberry, mosaic, 647
recovery from and acquired tolerance to, 743
Solanum, 168
Sour cherry, yellow leaf, 821
sugar beet, curly top, 825
sugar cane, Fiji disease, 512
systemic brooming of various trees, 8
tobacco, curly top, 743, 828
mosaics, 1, 2, 7, 17, 20, 22, 242, 250, 321, 441, 495, 684
virus-1, 16
necrosis, 20, 83, 323
thermal inactivation rates compared, 20
yellow ring spot, 549
tomato, mosaic, 757
tip blight (new), 156
of trees, 8
tulip "breaking," 3, 1001
turnip, mosaic, 323
Virus-like diseases, of trees, 8
Vitis californica, Plasmopara viticola overwintering in, 822
Volatile compounds, toxicity, testing method, 641

WADLEIGH, C. H., (756)
WAITE, M. B., (380)
WALKER, J. C., 120, 188, (641)
WALLACE, JAMES M., 743, 828
Walnut, black, virus-induced brooming, 8
Persian, bacteriosis, control by insoluble copper sprays, 826
WALTER, JAMES M., 23, 551
Water, injury to fruit of kumquat by, 214
sea spray, effect on cypress canker fungus, 823
Watermelon, blossom-end rot due to Pythium spp., 391
Pythium root and fruit rots of, 624
root knot, chemical soil treatment against, 25
wilt-resistant Hawkesbury variety, 5
Water-soaking, of plant tissues, effect on bacterial infection, 2
factors influencing, 2
tobacco leaves injured by, 884
WEAN, ROBERT E., 895
Web blight, of beans, Corticium microsclerotia causing, 559
WEBER, GEORGE F., 559
Weeds, chemical soil treatment against, 25
as reservoirs of Granville tobacco wilt, 22
root knot incidence and control on, 752
seeds killed by electrical pasteurization of soil, 18
WEETMAN, L. M., (21)
Weevils, chlorine gas seed disinfection for, 914
WEIMER, J. L., (989)
WEINDLING, R., (1), 23, 755
WEISS, FREEMAN, 23, 378, 1002
WELCH, D. S., 23
WELLMAN, FREDERICK L., 24, (757), 945
Wheat, bunt, chlorine gas seed disinfection for, 913
reinoculation of resistant varieties with purified physiologic races, 863
generalized standard errors in tests, review, 657
chlorotic leaf mottling, Tilletia levis causing, 456
Early Blackhull variety, crosses resistant to stem rust, 644
Fusarium spp. in diseases of, 7
leaf rust, inoculum source for fall infection, 4
physiologic races in 1938 epidemic, 962
resistant hybrids, 965
loose smut, infection factors, 717
seed treatments, for seedling blight and root rot, 16

stem rust, control by toluene compounds, 978
Early Blackhull variety resistant to, 644
epidemiology in Mexico, 22
as influenced by barberry eradication, 24
physiologic races, in Mexico, 22
temperature effects on infection and development of physiologic races, 4
WHITE, D. P., (367)
White-pine blister rust (See Cronartium ribicola)
White root rot, apple, 5
White rust, spinach, 12
White streak, narcissus disease (new) virus-induced, 826, 890
WIANT, JAMES S., 616
WILCOX, MARGUERITE S., 136
WILCOXON, FRANK, (16)
WILDE, S. A., 367
Wildfire, tobacco, as influenced by water-soaking of leaves, 884
WILSON, E. E., 828
Wilt, corn bacterial, as influenced by nitrogen supply, 17, 1051
motile bacterium in relation to, 1062
new host, 6, 284
cotton, infestation with nematodes plus, varietal reactions to, 755
nematodes in relation to, 752
cotton Fusarium-, as influenced by: acidity and nitrogen source and level, 756
nitrogen, phosphorus and potassium nutrition, 756, 757
unbalanced fertilization, 756
inoculation technique, 755
pathogenicity differences in strains, 1
soil contamination exp. technique, 756
varietal susceptibility to root-knot nematode plus, 974
water-culture infection method, 23
cotton Verticillium-, control and breeding for resistance, 753
cowpea, breeding for resistance, 826
cucumber bacterial, resistant hybrids, 996
lespedeza bacterial, 15
pepper Verticillium-, 359
potato Lygus-, non-identity with purple-top, 14
tobacco Granville-, control by rotation, 22
weeds as reservoirs of, 22
tomato Fusarium-, chemical soil treatment for, 25
cultural characters and pathogenicity of strains, 24
host resistance and pathogenicity, study technique, 945
hybrid resistant to, 757
varietal resistance, 25
and factors influencing, 871
watermelon, variety resistant to, 5
Wineberry, Japanese, Corticium galactinum infecting, 5
WINGARD, S. A., 9, (9), (18)
Witches' broom, virus-induced, in China, 460
WOLF, FREDERICK A., 103, (177), 194, 291, 915
WOOD, F. C., 728
Wood, decay by Ustulina and other Ascomycetes, 2
fungi destroying and staining, relative importance and seasonal prevalence in U. S. A., 1031
gum-pocket formation in sweet cherry, microchemistry of, 20
preservatives, acceleration of toximetric tests by soil, 901
WORMALD, H., 915
Wound dressings, effect on apple perennial canker, 739
WRIGHT, ERNEST, (14), (485)
WRIGHT, L. K., 24
Writing, brevity in, 292
WYCKOFF, RALPH W. G., (83)

Xanthium americanum, Corticium microsclerotia on, 561
X-ray diffraction, of tobacco mosaic virus proteins prepared by Na_2SO_4 method, 17
X-virus, potato, on American vs. European varieties, 57
immunity inheritance, 362
inhibition by insect juices, 323
Xylaria, hardwood decay by spp., 2

YAP, FRANCIS, (596)
YARWOOD, C. E., 24, 282, 288, 828, 933
Yeasts, pines infected by, 1036
Yellow cuprous oxide, as fungicide of small particle size, 9
Yellow dwarf, potato, virus inhibition by insect juices, 2, 321, 322, 323
Yellow leaf, sour cherry, virus(?)-induced, 821
Yellow mosaics, potato, comparisons and classification, 917
Yellow ring spot, tobacco, symptoms, 549
virus "longevity" in seed, 549
Yellow spot, celery, aphid transmission, 824
pineapple, geographical distribution and insect vector, 285
Yellows, aster, in delphinium, 826
relation to potato purple-top, 14
YORK, HARLAN H., 24
YOTHERS, M. A., (739)
YOUNG, J. E., (895)
YOUNG, P. A., 25, 871
YOUNG, V. H., 756
Youngberry, stamen blight due to Halalosphaeria deformans, 829
Y-virus, potato, on American vs. European varieties, 51
YU, T. F., 448, 459

ZAUMEYER, W. J., 9, 25
ZELLER, S. M., 829